



## Critical review on *Agnimantha* (*Premna integrifolia* Linn and *Clerodendrum phlomidis* Linn)

**Dr. Umakant N Rabb**

Associate Professor, Department of Dravyaguna Vijnana, Shri Ravi Patil Health and Education Society's, Dr. Ravi Patil Ayurvedic Medical College, Hospital and Research Centre, Honaga, Belagavi, Karnataka, India

\* Corresponding Author: **Dr. Umakant N Rabb**

---

---

### Article Info

**ISSN (online):** 2582-7138

**Volume:** 03

**Issue:** 04

**July-August** 2022

**Received:** 03-07-2022;

**Accepted:** 20-07-2022

**Page No:** 292-298

### Abstract

The *Agnimantha* i.e *Premna serratifolia* Linn. Synonym *Premna integrifolia* Linn is a large, thorny deciduous shrub or a tree, up to 9 meters in height, commonly seen at Indian and the Andamans coasts. The Agni means fire and *Mantha* means churning. When the plant woods rubbed each other, generate fire, so named as *Agnimantha*. In *Ayurveda* the roots and leaves are most used parts for various medicinal preparations. The roots are considered as astringent, stimulant, liver tonic, laxative, carminative and antibacterial. Leaves infusion with pepper is useful in cold and fever, leaf sap is useful to alleviate headache. It can be used as hypoglycemic, anthelmintic, aphrodisiac, anti-inflammatory, analgesic and antimicrobial agent. It cure diseases like urticaria, asthma, haemorrhoids, erysipelas, fever, worms, allergic conditions. It is described in one of the ingredient in Dashamoola. The *Agnimatha* is of two types Viz; *Laghu Agnimatha*(*Premna integrifolia* Linn), and *Brihat Agnimantha*(*Clerodendrum phlomidis* Linn). The present review dealt with the brief introduction and pharmacological actions, its traditional uses, modern research activities and also discussed the botanical description, chemical composition, biological activities on the basis of digital scientific database

**Keywords:** *Agnimantha*, *Premna integrifolia* Linn, *Premna mucranata*, *Clerodendrum Phlomidis* Linn, Pharmacological actions

---

---

### Introduction

The *Premna serratifolia* Linn. (Syn. *Premna integrifolia* Linn), is a large, thorny deciduous shrub or a tree, up to 9 meters in height, commonly seen at Indian and the Andaman coasts. In *Ayurveda* the roots and leaves are most used parts for various medicinal preparations. The roots are considered as astringent, stimulant, liver tonic, laxative, carminative and antibacterial. Leaves infusion with pepper is useful in cold and fever, leaf sap is useful to alleviate headache<sup>[1, 2]</sup>. The roots are useful in vitiated conditions of *Vata*, *Kapha*, neuralgia, inflammations, cardiac disorders, cough, asthma, bronchitis, leprosy, skin disorders, dyspepsia, flatulence, constipation, fever, diabetes and anorexia<sup>[3]</sup>. In Vietnamese traditional medicine, the leaves of *Schisandra Chinensis* (Turcz) Ball belongs to Schisandraceae family, are used to treat indigestion, dysuria and dysentery; roots against indigestion, stomach ache and fever<sup>[4]</sup>. The root of methanolic extract showed better immunomodulatory activity in experimental mice. A paste of its roots with water and butter fat is given in urticaria.

### Discussion

Various pharmacological actions along with its formulations discussed systematically;

#### *Rasa Panchaka* <sup>[5]</sup>

**Rasa:** *Tikta* (Bitter), *Katu* (Pungent), *Kashaya* (Astringent)

**Guna:** *Ruksha* (Dryness), *Laghu* (Light for digestion)

**Virya:** *Ushna* (Hot in potency)

**Vipaka:** *Katu* (Pungent)

**Karma:** *Shothahara* (Anti-inflammatory), *Vedanasthapana* (Which relieves pain)

**Doshagnata:** *Kapha-Vatahara* (Subside *Kapha* and *Vata* Doshas)

### Vishistha Yogas

*Amritarishta, Dantadyarishta, Dasamularishta, Agastya Haritaki Rasayana, Chyavanaprasha, Brahma Rasayana, Dashamula Katutraya Kwatha Churna, Rasnadi Kwatha Churna, Saptasara Kwatha Churna, Indukanta Ghrita, Dashamulaghrita, Dhanvantara Ghrita, Mahapanacagavya Ghrita, Lasunadi Ghrita, Sukumara Ghrita, Dhanvantara Taila, Narayana Taila, Mahanarayana Taila, Mahavishagarbha Taila, Sahacharadi Taila*<sup>[6]</sup>. *Guduchyadi Taila, Dashamula Taila, Vajrakapata Rasa, Shirahsuladivajra Rasa*<sup>[7]</sup>, *Dashamularka, Tagaradi Kwatha Churna, Bhunimbadi Astadashanga Kwatha Churna, Viratarvadigana Kwatha Churna*<sup>[8]</sup>.

### Botanical Description<sup>[9]</sup>

It is small tree, branches are spiny, bark is thin pale and exfoliating wood is light brown and scented. The leaves are ovate or ovate oblong, long acuminate base rounded cordate or tapering, entire or irregularly dentate blade 7-15 cms long petiole is 2.5 cms long.

The flowers are arranged in terminal, corymbose, trichotomous panicles and are greenish in colour, calyx comprised of 4-5 sepals, with rounded and nearly equal teeth. Corolla lobes are equal or bilabiate, upper lip retuse or emarginate, lower lip of 3 equal lobes and throat closed with white hair. Fruit is globose drupe, green when young, dark at maturity, 3-5mm in diameter. The mature trees start flowering in April and fruiting in May and June.

### Chemical Composition

Premnacorymboside A, scutellarioside II, quercetin - 3-rutinoside, leonurioside A were isolated from the stem bark<sup>[10]</sup>. Alkaloid contents like premnine, ganikarine, premnazole are reported in root from different researchers<sup>[11]</sup>. 1 $\beta$ , 3 $\alpha$ , 8 $\beta$ -trihydroxy-pimara-15-ene; 6 $\alpha$ , 11, 12, 16- tetrahydroxy-7-oxo-abieta-8, 1, 13-triene; 2 $\alpha$ , 19- dihydroxypri -mara-7,15-diene were identified in root bark<sup>[12]</sup>.

### Synonyms<sup>[13]</sup>

*Arani, Ketu, Tarkari, Vaijayantika, Mathana, Vanhimathana, Ganika, Ganikarika, Pavana, Jayanti, Vijaya, Jaya.*

### Gana

According to *Acharya Charaka*<sup>[14]</sup> *Shothahara, Sheetaprashamana, Anulomana.*

According to *Acharya Sushruta*<sup>[15]</sup> *Brihatpanchamula, Vatasanshamana, Viratarvadi, Varunadi*

According to *Acharya Vagbhata*<sup>[16]</sup> - *Brihat Panchamula,*

According to *Dhanwantari Nighantu*<sup>[17]</sup> *Guduchyadi Varga*

According to *Raja Nighantu*<sup>[18]</sup> *Prabhadradi Varga*

According to *Madanapala Nighantu*<sup>[19]</sup> *Abhayadi Varga*

According to *Kayadeva Nighantu*<sup>[20]</sup> *Oshadhi Varga*

According to *Priya Nighantu*<sup>[21]</sup> *Haritakyadi Varga, Brihat Panchamula*

According to *Bhavaprakasha Nighantu*<sup>[22]</sup> *Guduchyadi Varga*

According to *Chandra Nighantu*<sup>[23]</sup> *Niruha Gana*

### Pharmacological actions as per Ayurvedic science<sup>[24]</sup>

**Dosha Karma-** Due to *Ushna Virya* it subsides *Kapha* and *Vata* Doshas. Therefore it is recommended in *Kapha Vatajanya* diseases

**Sansthanika Karma (Systemic actions)-Bahya:** External Application- As it is *Ushna* it alleviates *Shotha* (Edema), and *Vedana*(Pain). Therefore it is used in edema and pain.

**Pachana Sansthana:** Due to *Ushna Virya* it acts as *Dipana* (Appetizer), *Pachana* (Digestive), and *Anulomana*. Therefore it is indicated in *Agnimandya* (Loss of appetite), *Amadosha, Vibandha* (Constipation).

**Raktavaha Sansthana:** The *Agnimatha* is *Raktashodhaka* (Purifies the blood), *Hridayottejaka* (Cardiac stimulant), and *Shothahara* (Subsides edema). Therefore it is advised in *Raktavikara* (Blood disorders), *Hridaya Daurbalya* (Strengthen the cardiac muscle), and edema.

**Shwasana Sansthana:** As it is *Kaphaghna* used in *Kaphaja* diseases.

**Mutravaha Sansthana:** It acts as *Pramehaghna* (Urinary disorders including diabetes) used in *Prameha Roga* (Urinary diseases including diabetes).

**Tapakrama:** It is *Jwaraghna* (Anti pyretic) therefore used in *Jwara* (Fever).

**Satmikarana:** As it is *Katu Pausthika*, it is advised in post fever general debility, *Pandu* (Anemia).

**Part used:** *Patra Swarasa* (Leaves juice), *Twak* (Bark), *Moola* (Root), *Moola Twak* (Root bark).

**Dosage:** Leaf juice 10-20ml, Churna (Powder)-1-3 grams, *Kwatha* (Decoction)-50-100 ml.

**Bheda-(Types):** *Kshudra Agnimantha* and *Brihat Agnimatha*

### Pharmacological actions according to modern

- **Analgesic Activity:** Analgesic activity was evaluated by using ethanolic extract of leaves of *P. integrifolia*<sup>[25]</sup>. Analgesic activity was also evaluated using methanolic extract of *Premna integrifolia* (MEPI) bark by writhing test in rats at doses 100 and 200 mg/kg body weight<sup>[26]</sup>
- **Anti-arthritic activity:** Anti-arthritic activity was evaluated using ethanol extract of *P. serratifolia* wood by Freund's adjuvant induced arthritis model<sup>[27]</sup>
- **Anti-bacterial activity:** Antibacterial activity of ethanolic extract of leaves of *Premna integrifolia* was tested by using the disc diffusion method. The extract showed significant antibacterial activity against both gram positive and gram negative bacteria. Alcoholic extract of the root bark of *P. integrifolia* showed good antibacterial activity against gram-positive organisms during preliminary screening<sup>[28, 29]</sup>
- **Anti-cancer/ anti-tumour/ cytotoxicity/tumour suppression activity:** Anticancer activity of biosynthesized silver nanoparticles (AgNps) using the ethanolic leaf extract of *P. serratifolia* was evaluated against carbon tetra chloride (CCl4) induced liver cancer in Swiss albino mice (Balb/c)<sup>[31, 32]</sup> The methanolic extracts of leaves, root barks (RB) and root wood of *P. serratifolia* for cytotoxic activity against two cancer cell lines<sup>[33]</sup>. The alcoholic extract of leaves of *P. serratifolia* possessed significant antitumor activity<sup>[34]</sup>. The cytotoxicity screening system validates anticancer use of the plant by traditional healers and literature claims<sup>[35]</sup>.
- **Anti-inflammatory activity:** Anti-inflammatory activity of methanolic extract of *Premna integrifolia* (MEPI) bark was evaluated using carrageenan induced

- inflammatory model in rats at doses 100 and 200 mg/kg b. w [37]. The administration of the root extract inhibited the carrageenan induced acute paw oedema and formalin induced chronic paw oedema in a dose dependent manner [38]. The root and root callus extracts of *P. serratifolia* augments its anti-inflammatory activity against carrageenan induced paw edema. The results of a study proved the anti-inflammatory effect of callus derived luteolin through its biologically active components, which may worth for further investigation and elucidation [39]. Anti-inflammatory activity was evaluated by pretreatment with a single dose of methanolic extract of *P. integrifolia* (PIM) (300 mg/kg b.w.). The extract also showed significant inhibition of cyclo-oxygenase (COX-I) activity on rat uterus and plasma membrane stabilization. The results scientifically demonstrated the anti-inflammatory activity of *P. integrifolia* roots in various experimental models probably through their antihistaminic, antikinin, COX-inhibitory and antioxidant action [40]
- **Anti-microbial activity-** Antimicrobial activity was evaluated against the selected human pathogens (*Bacillus* sp., *Enterococcus faecalis*, *E. coli*, *Klebsiella pneumoniae*, Non-haemolytic *Streptococci*, *Streptococcus epidermidis*, *Pseudomonas aeruginosa*, *Salmonella typhimurium*) by using natural leaves, roots and its respective calli induced with help of various plant growth regulators. Results revealed increased inhibitory activities of callus extracts. The activities were found to be better when compared to the natural plant material extracts [41]. *P. serratifolia* was screened to evaluate in-vitro antimicrobial activity against the selected human pathogenic organisms [42]. The antimicrobial activity of different extracts of bark and wood of *P. serratifolia* was also studied against nine bacterial and four fungal organisms by disc diffusion method [43].
  - **Anti-obesity or hypolipidemic activity:** Anti-obesity activity of chloroform-methanol (1:1) extract of *P. integrifolia* (CMPI) in mice fed with cafeteria diet was evaluated [44]. The effect of aqueous enriched fraction of *P. integrifolia* root (AEFPIR) against cafeteria diet induced obesity in Swiss Albino Mice was studied. There was a significant decrease in body weight, BMI, food consumption and in the levels of serum glucose, triglyceride, total cholesterol, LDL, and VLDL with a significant increase in the level of HDL in mice treated with simvastatin and AEFPIR groups compared with cafeteria diet group [45]. Hypolipidemic activity was also evaluated using hypercholesterolemic rabbit model in comparison to the established hypolipidemic drug atorvastatin [46]. Anti-hyperlipidemic activity of *P. integrifolia* was evaluated in an experimental animal model. High fat diet caused significant increase in serum cholesterol, triglyceride, VLDL, LDL and significant reduction in HDL level. Both *P. integrifolia* and atorvastatin treatment showed significant prevention in increased in serum cholesterol, triglyceride, LDL as compared to cholesterol control group. HDL level was significantly increased in both treated and standard control group as compared to cholesterol control group [47]. Anti-hyperlipidaemic activity of *P. integrifolia* on nicotine induced hyperlipidaemia was evaluated. From the above results, it is clear that *P. integrifolia* is an effective anti-hyperlipidemic agent [48].
  - **Anti-oxidant activity:** Ethanolic extract of root and root derived callus extracts of *P. serratifolia* were investigated against paracetamol induced oxidative stress in blood of male albino rats. This investigation concluded that, ethanolic extract of root callus possess significant antioxidant activity, which can help to overcome paracetamol induced toxicity [49]. *P. serratifolia* possessed radical scavenging activity [50]. The free radical scavenging activity of *P. serratifolia* leaf has been evaluated in various in-vitro model systems. The study showed that the methanolic extract of plant has free radical scavenging activity against superoxide radical, nitric oxide radical, hydroxyl radical, DPPH radical, ABTS radical and inhibition of lipid peroxidation [51]. The isolated compound, 11,12,16-trihydroxy-2-oxo-5-methyl-10-demethyl-abieta-1,10,6,8,11,13-pentene that appears to be a novel compound based on a new diterpene skeleton also possesses in-vitro antioxidant effects [52]. Antioxidant properties of the plants were analyzed by determining the scavenging effects of free radicals such as superoxide, hydroxyl, nitric oxide and lipid peroxidation generated with in-vitro assay systems [53]. Methanolic extract of *P. integrifolia* showed significant anti-oxidant activity [54]. Antioxidant activity was evaluated by using methanolic extract of *P. serratifolia* leaf in paracetamol intoxicated rats. The study revealed that the methanolic extract of *P. serratifolia* was found to have potential antioxidant activity in the animal model system [55]. Antioxidant effect was monitored using the DPPH radical scavenging assay. Through the use of comprehensive spectroscopy studies, the isolated active antioxidant principle was identified as acteoside (verbacoside) which is about four times more active than the crude root wood extract [56]. Significant scavenging activity was observed in DPPH and NO assays. IC50 values of compounds 10-O-trans-p-coumaroylcatalpol and 4-hydroxy-E-globularinin were 0.37 and 0.29 M/mL in DPPH free radical scavenging assay, respectively [57]. Antioxidant activity of the oil and various organic extracts of *P. integrifolia* leaves were examined using DPPH and superoxide radical scavenging activities assay. The amount of total phenolic compounds was also determined as gallic acid equivalent. The results of this study suggest that the natural products derived from *P. integrifolia* may have potential use in food, pharmaceutical and/or cosmetic industries [58]. The extract showed significant antioxidant activity which is scientifically demonstrated the use of *P. integrifolia* as a potential source of natural antioxidant [59]. The extracts of *P. integrifolia* root possesses beneficial effect on human leucocytes and erythrocytes against H<sub>2</sub>O<sub>2</sub> induced oxidative damage which substantiates their use in ethnomedicine as an antioxidant. Observed effect can be attributed to the flavonoid and phenol contents in the plant [60].
  - **Anti-parasitic activity:** Antiparasitic activity of a few New Caledonian medicinal plants were studied [61]. Among the selected plants, *Scaevola balansae* and *P. serratifolia* were the most active against *Leishmania donovani* with IC<sub>50</sub> values between 5 and 10 g/ml
  - **Anti-ulcer and gastroprotective activity:** Antiulcer activity of ethanolic extract of leaves of *P. serratifolia* using aspirin induced ulcers at dose levels of 200 and 400 mg/kg compared with ranitidine as a

standard drug. Results showed dose dependent decrease in ulcer index, gastric acid secretion, free acidity, total acidity and increased the pH of gastric juice in aspirin and extract treated group of animals. Histopathological studies revealed that high doses of *P. serratifolia* leaves prevented ulcer formation [62].

- **Cardiac stimulant or cardioprotective activity:** Cardiac stimulant activity of stem bark and stem-wood of *P. integrifolia* was evaluated by using Isolated Frog Heart Perfusion Technique [63]. Cardioprotective effect of ethanol extract of stem-bark and stem-wood of *P. serratifolia* were tested on isoproterenol induced experimental myocardial infarction in rats. The cardioprotective effect of ethanol extract of stem-bark and stem-wood of *P. serratifolia* on isoproterenol induced myocardial infarction in rats. The protective myocardial effect may be due to the phytoconstituents such as iridoid glycosides, alkaloids, flavonoids and phenolic compounds [64].
- **CNS depressant activity:** CNS depressant activity of methanolic extract of *P. integrifolia* (MEPI) bark was evaluated using hole cross and open field test in rats at doses 250 and 500 mg/kg b. w. The depressing effect in hole-cross test was most intense during the second (60 min) and third (90 min) observation periods [65].
- **Hepatoprotective activity:** Hepatoprotective activity of alcoholic extract of leaves of *P. serratifolia* using carbon tetrachloride was studied. The degree of protection in hepatoprotective activity was measured by using biochemical parameters such as SGOT, SGPT, ALP, bilirubin and total protein. The results suggest that the alcoholic extract at the dose level of 250 mg/kg produced significant hepatoprotection by decreasing the activity of serum enzymes, bilirubin, and lipid peroxidation which is comparable to that of standard drug silymarine [66]. Hepatotoxicity induced by paracetamol (800 mg/kg) administration was studied. The effects of root callus extracts were comparable to that of standard drug, silymarine. Histopathological findings also suggested the root callus extracts of *P. serratifolia* preventing the development of chronic damage [67].
- **Hypoglycemic activity:** Hypoglycemic activity was evaluated using streptozotocin induced type 1 and type 2 diabetic rats [68]. Hypoglycemic activity was also evaluated using 95% ethanolic extract of leaves in alloxan induced diabetic rats. Steady diabetes was confirmed by noting urine sugar regularly and then measuring blood glucose values before starting the experiment. To confirm definite hypoglycemic effect, final blood glucose values of each sample were brought down closer to normal fasting level using different doses of the ethanolic extract. The results indicate that *P. integrifolia* showed definite blood glucose lowering effect within 1 week using 250 mg/kg double dose [69].
- **Immunomodulatory activity:** Immunomodulatory activity was evaluated by using methanol extract of root of *P. integrifolia* in BALB/c mice [70].
- **Anti-ageing or life promoting or anti-oxidant activity:** In modern times, there has been an increase in the use of plants or herbal constituents for the prevention of age-related disorders. 4-Hydroxy-E-globularinin (4-HEG) is an iridoid and a major component of *P. integrifolia*. The discovery of this component represents a break-through in geriatrics because of its longevity-

promoting activity of 4-HEG in an animal model. 4-HEG may serve as a lead compound for the development of important nutraceuticals preventing the aging process [71]. Aging, the major cause of several ailments has led to intense exploration of potential drugs that delay aging and its associated effects [72].

- **Neuroprotective activity:** Neuroprotective effects were evaluated by using roots of *P. serratifolia* in the experimental model of febrile seizure [73].

### Aims and Objectives

- To provide brief introduction and pharmacological actions, its traditional uses, modern research activities and also discussed the botanical description, chemical composition, biological activities on the basis of digital scientific database
- To provide comprehensive information on the pharmacological activities of various parts of *Premna integrifolia* Linn. All the relevant universally accepted electronic databases were searched with respect to the terms *Agnimantha*, *Premna integrifolia* Linn, *Premna obtusifolia* Linn, *Premna serratifolia* Linn including Indian classical texts, pharmacopoeias, journals.

### Material and Methods

- A survey was done by online as well as *Ayurvedic* classical texts books and the data were obtained by several electronic scientific databases and the additional information is obtained from the various *Ayurvedic* texts.

### Conclusion

The present review reveals *P. integrifolia* to possess analgesic/antinociceptive, anti-arthritic, antibacterial, anticancer/antitumor/cytotoxic/tumor suppression, anti-inflammatory, antimicrobial, anti-obesity/hypolipidemic, antioxidant, antiparasitic, antiulcer/gastro-protective, cardiac stimulant/cardioprotective, CNS depressant, hepatoprotective, hypoglycemic, immunomodulatory, longevity-promoting and neuroprotective activities. According to the literature, most of the pharmacological activities of *P. integrifolia* is investigated by using methanol and ethanol as extractive solvents. The activities may be due to the presence of these phytochemicals in the extracts. The present information will be useful for setting up of research protocols for modern drug and *Ayurvedic* formulation development in curing and treating various diseases, which can prove its efficacy as a novel source for new drug development. The information collected here will be useful to set-up research protocols for modern drugs and *Ayurvedic* formulation development.

### References

1. Kirtikar KR, Basu BD. Indian Medicinal Plants. Lalith Mohan Basu, 49-Leader Road, Allahabad, India, Edition. 1994; 2(3):1927-1929.
2. Kapoor LD. CRC Hand book of Ayurvedic Medicinal Plants- Herbal Reference Library, CRC press, Boca Raton, London, Newyork, Washington DC, 2001, 70.
3. George KV, Samuel KA, Abraham J, George BP. Investigations on Bioactive compounds of *Premna integrifolia* Linn. *Aryavaidyan*. 2003; 16(4):234-239.
4. Sharma PV, Dravyaguna Vijnana, Chaukambha Bharati Academy, Varanasi, Edition Reprint. 2009; 2:221.
5. Sharma PV, Dravyaguna Vijnana, Chaukambha Bharati

- Academy, Varanasi, Edition Reprint. 2001; 2:222.
6. Government of India, Ministry of Health and Family Welfare, Department of Indian Systems of Medicine and Homoeopathy; 1<sup>st</sup> edition, New Delhi Formulary Single Drugs, List of single drugs of plant origin. The Ayurvedic Formulary of India, 1978, 1:241-59.
  7. Government of India, Ministry of Health and Family Welfare Department of Indian Systems of Medicine and Homoeopathy. 2<sup>nd</sup> edition New Delhi. Formulary Single Drugs, List of single drugs of plant origin. The Ayurvedic Formulary of India. 2000; 1:307-330.
  8. Government of India, Ministry of Health and Family Welfare Department of Indian Systems of Medicine and Homoeopathy. 1<sup>st</sup> edition, New Delhi, Formulary Single Drugs, List of single drugs of plant origin. The Ayurvedic Formulary of India. 2000; 2:327-340.
  9. <https://vikaspedia.in>
  10. Hang NTB, Ky PT, Minh CV, Cuong NX, Thao NP, Kiem PV. Study on the Chemical Constituents of *Premna integrifolia* Linn. Natural Product Communications. 2008; 3(9):1449-1452.
  11. Gokani RH, Lahiri SK, Santani DD, Shah MB. Evaluation of immunomodulatory activity of *C. phlomidis* and *Premna integrifolia* root. International Journal of Pharmacology. 2007; 3(4):352-356.
  12. Yadav D, Tiwari N, Gupta MM. Simultaneous quantification of diterpenoids in *Premna integrifolia* using a validated HPTLC method. J Sep. Sci. 2011; 34:286-91.
  13. Acharya Kaiyadeva, Kaiyadeva Nighantu, Oshadhi Varga, Shloka No- 47,48, redacted by Sharma P.V., 1st edition, New-Delhi, Chaukhambha publications, 1979, 8.
  14. Acharya Charaka, Charaka Samhita, Sutrasthana 4<sup>th</sup> Chapter, Shloka No-38, Shloka No-42, Shloka No-Charaka Chandrika Hindi commentary of Agnivesha, edited by Dr. Bramhananda Tripathi, Chaukhambha Surabharati Prakashana, Varanasi, Reprint, 2002, 81-83.
  15. Acharya Susruta, Sushruta Samhita, Sutrasthana 38<sup>th</sup> Chapter, Shloka No- 10, Shloka No-34, Shloka No-8, Uttarstthchapter, Edited with AyurvedaTattva-Sandipika, By Kaviraj Ambikadutta Shastri, Varanashi, Chaukhamba Sanskrit Sansthan, Reprint, 2014, 183-188.
  16. Acharya Vagbhata. Astanga Hridayam of Srimadvagbhata, 10<sup>th</sup> Chapter, 10-19, Shloka No-167, Edited with Nirmala Hindi commentary, By Dr. Bramhanand Tripathi, Delhi, Chaukhamba Sanskrit Pratishthan, Reprint, 2014, 120.
  17. Prof Priya Vrat Sharma, Dr. Guru Prasad Sharma. Dhanwantari Nighantu, Shloka No- 108,109, Guduchyadi Varga, Varanasi, Chaukhmbha Orientalia, 3rd Edition 2002, 35.
  18. Pandit Narahari, Raj Nighantu, Prabhadradi Varga, Shloka No- 22-25, redacted by Indradeva Tripathi, Varanasi, Chaukhamba Krishnadas Academy, 2003, 268.
  19. Bhagwan Dash, Madanapal Nighantu, Abhayadi Varga, Shloka No- 47, B. Jain Publisher's pvt. Ltd, first edition, New Delhi, 1991, 15.
  20. Acharya Kaiyadeva, Kaiyadeva Nighantu, Oshadhi Varga, Shloka No- 47,48, redacted by Sharma P.V., 1st edition, New-Delhi, Chaukhambha publications, 1979, 8.
  21. Prof. Priya Vrat Sharma, Priya nighantu, Haritakyadi Varga, Shloka No- 28-31, Hindi commentary entitled Padma, Varanasi, Chaukhambha Surabharati Prakashana Edition 2004, 10-11.
  22. Acharya Bhavamishra. of Bhavaprakasha Nighantu, Guduchyadi Varga, Shloka No-23,24, by Dr. Bulusu Sitaram, Volume 1, Chaukhambha Orientalia, Varanasi, First edition, 2006, 234.
  23. Kashmiri Vidyaraj Chandranandana, Chandra Nighantu (Madanadi Nighantu) Niruha Gana, Shloka No-8-10, edited by Acharya Balakrishna, Haridwar, Divya Prakashana, First edition, 2015, 42.
  24. Sharma PV Dravyaguna Vijnana, Chaukhambha Bharati Academy, Varanasi, Reprint. 2001; 2:221-223.
  25. Kumar KU, Soma P, Kumar SS, Chandra SM, Kumar BS. Assessment of analgesic and antibacterial activity of *Premna integrifolia* Linn. Leaves. Int J Pharm Sci., Res. 2011; 2:1430-5.
  26. Khatun H, Majumder R, Mamun A, Alam EK, Jami SI, Alam B. Preliminary pharmacological activity of the methanolic extract of *Premna integrifolia* barks in rats. *Avicenna J Phytomed.* 2014; 4:215-24.
  27. Rajendran R, Krishnakumar E. Anti-arthritis activity of *Premna serratifolia* Linn. Wood against adjuvant induced arthritis. *Avicenna J Med Biotech.* 2010; 2:101-106.
  28. Kurup KK, Kurup PA. Antibiotics substances from the root bark of *P. integrifolia*. *Die Naturwissenschaften.* 1964; 20:484.
  29. Rahman A, Shanta ZS, Rashid MA, Parvin T, Afrin S, Khatun MK, *et al.* *In-vitro* antibacterial properties of essential oil and organic extracts of *Premna integrifolia* Linn. *Arabian J Chemist*, 2011.
  30. Paul John JA, Selvi BK, Karmegam N. Biosynthesis of silver nanoparticles from *Premna serratifolia* L. leaf and its anticancer activity in CCl<sub>4</sub>-induced hepato-cancerous Swiss albino mice. *Appl Nanosci*, 2015, 1.
  31. Sridharan G, Brindha P, Jaiganesh C, Sivasubramanian S. Anti-tumor potential of *Premna integrifolia* Linn. against Ehrlich ascites carcinoma cell lines. *Pharmacologyonline.* 2011; 2:438-450.
  32. Selvam TN, Venkatakrishnan V, Damodar KS, Elumalai P. Antioxidant and tumor cell suppression potential of *Premna serratifolia* Linn leaf. *Toxicol Int.* 2012; 19:31-34.
  33. Habtemariam S, Varghese GK. A novel diterpene skeleton: Identification of a highly aromatic, cytotoxic and antioxidant 5-Methyl-10-demethyl-abietane-type diterpene from *Premna serratifolia*. *Phytother Res.* 2015; 29:80-5.
  34. Vadivu R, Jerad S, Girinath K, Kannan PB, Vimala R, Kumar NM. Evaluation of hepatoprotective and in-vitro cytotoxic activity of leaves of *Premna serratifolia* Linn. *J Sci Res.* 2009; 1:145-152.
  35. Biradi M, Hullatti K. Screening of Indian medicinal plants for cytotoxic activity by Brine Shrimp Lethality (BSL) assay and evaluation of their total phenolic content. *Drug Dev Ther.* 2014; 5:139-144.
  36. Khatun H, Majumder R, Mamun A, Alam EK, Jami SI, Alam B. Preliminary pharmacological activity of the methanolic extract of *Premna integrifolia* barks in rats. *Avicenna J Phytomed.* 2014; 4:215-224.
  37. Rathore RS, Prakash A, Singh PP. Preliminary study of anti-inflammatory and anti-arthritis activity. *Rheumatism.* 1997; 12:130.

38. Sasidharan N Peechi. Kerala Forest Research Institute; Qualitative and quantitative analysis of biologically active principles, baicalein, luteolin and psoralen from *Oroxylum indicum*, *Premna serratifolia*, *Aegle marmelos* and their Allied Species. KFRI Research Report No. 350 (KFRI/486/05), 2010, 1-57.
39. Singh CR, Nelson P, Boopathy S. *In-vitro* conservation and protective effect of *Premna serratifolia* Linn. An important medicinal tree. *Int J Pharm Appl*. 2012; 3:332-343.
40. Gokani RH, Lahiri SK, Santani DD, Shah MB. Evaluation of anti-inflammatory and antioxidant activity of *Premna integrifolia* root. *J Complement Integr Med*, 2011.
41. Singh CR. Antimicrobial effect of callus and natural plant extracts of *Premna serratifolia* L. *Int J Pharm Biomed Res*. 2011; 2:17-20.
42. Rajendran R, Basha NS. Antimicrobial activity of crude extracts and fractions of *Premna serratifolia* Linn. root. *Medicinal Plants*. 2010; 2:33-38.
43. Rajendran R. Antimicrobial activity of different bark and wood of *Premna serratifolia* Linn. *Int J Pharma Biosci*. 2010; 6:1-9.
44. Mali PY, Bigoniya P, Panchal SS, Muchhandi IS. Anti-obesity activity of chloroform-methanol extract of *Premna integrifolia* in mice fed with cafeteria diet. *J Pharm Bioallied Sci*. 2013; 5:229-236.
45. Mali PY. Effect of aqueous enriched fraction of *Premna integrifolia* root against cafeteria diet induced obesity in Swiss albino mice. *Int J Green Pharm*. 2013; 7:315-321.
46. Bagchi C, Tripathi SK, Hazra A, Bhattacharya D. Evaluation of hypolipidemic activity of *Premna integrifolia* Linn. Bark in rabbit model. *Pharmbit*. 2008; 18:149-153.
47. Patel MJ, Patel JK. Evaluation of the anti-hyperlipidemic activity of *Premna integrifolia* using experimental animal model. *Int J Res Phytochem Pharmacol*. 2011; 1:146-159.
48. Patel MJ, Patel JK. Evaluation of the anti-hyperlipidaemic activity of *Premna integrifolia* on nicotine induced hyperlipidaemia in rats. *Int J Pharma Biosci*. 2012; 3:226- 232.
49. Kumar KU, Soma P, Kumar SS, Chandra SM, Kumar BS. Assessment of analgesic and antibacterial activity of *Premna integrifolia* Linn. leaves. *Int J Pharm Sci Res*. 2011; 2:1430-1435.
50. Paul John JA, Selvi BK, Karmegam N. Biosynthesis of silver nanoparticles from *Premna serratifolia* L. leaf and its anticancer activity in CCl<sub>4</sub>-induced hepato-cancerous Swiss albino mice. *Appl Nanosci*, 2015, 1-8.
51. Selvam TN, Venkatakrishnan V, Damodar KS, Elumalai P. Antioxidant and tumor cell suppression potential of *Premna serratifolia* Linn leaf. *Toxicol Int*. 2012; 19:31-34.
52. Habtemariam S, Varghese GK. A novel diterpene skeleton: Identification of a highly aromatic, cytotoxic and antioxidant 5-Methyl-10-demethyl-abietane-type diterpene from *Premna serratifolia*. *Phytother Res*. 2015; 29:80-85.
53. Sasidharan N. Peechi: Kerala Forest Research Institute; Qualitative and quantitative analysis of biologically active principles, baicalein, luteolin and psoralen from *Oroxylum indicum*, *Premna serratifolia*, *Aegle marmelos* and their Allied Species. KFRI Research Report No. 350 (KFRI/486/05), 2010. Page No-1-57.
54. Gokani RH, Lahiri SK, Santani DD, Shah MB. Evaluation of anti-inflammatory and antioxidant activity of *Premna integrifolia* root. *J Complement Integr Med*, 2011, 8.
55. Selvam NT, Vengatakrishnan V, Damodar KS, Murugesan S. Evaluation of tissue level antioxidant activity of *Premna serratifolia* leaf in paracetamol intoxicated Wistar albino rats. *Int J Pharmaceut Life Sci*. 2010; 1:86-90.
56. Bose LV, Varghese GK, Habtemariam S. Identification of acteoside as the active antioxidant principle of *Premna serratifolia* root wood tissues. *Phytopharmacology*. 2013; 4:228-236.
57. Yadav D, Masood N, Luqman S, Brindha P, Gupta MM. Antioxidant furofuran lignans from *Premna integrifolia*. *Ind Crops Prod*. 2013; 41:397-402.
58. Rahman A, Sattar MA. Hyderabad: Omics Group Conferences; 2011. Antioxidant Properties of Essential oil and Organic Extracts of *Premna integrifolia* Linn. International Conference and Exhibition on Pharmaceutical Biotechnology.
59. Gokani RH, Lahiri SK, Santani DD, Shah MB. Evaluation of *in-vitro* anti-oxidant activity of *Premna integrifolia* Linn. root. *Res J Pharmacog Phytochem*. 2010; 2:196-199.
60. Mali PY. Beneficial effect of extracts of *Premna integrifolia* root on human leucocytes and erythrocytes against hydrogen peroxide induced oxidative damage. *Chron Young Sci*. 2014; 5:53-58.
61. Desrivot J, Waikedre J, Cabalion P, Herrenknecht C, Bories C, Hocquemiller R, *et al*. Antiparasitic activity of some New Caledonian medicinal plants. *J Ethnopharmacol*. 2007; 112:7-12.
62. Jothi ET, Karthikeyan R, Suryalakshmi PV, Srinivasababu P. Gastroprotective potential of *Premna serratifolia* Linn. leaves against aspirin induced ulcer in albino rats. *Pharmacologyonline*. 2010; 3:189-198.
63. Rajendran R, Suseela L, Meenakshi SR, Saleem BN. Cardiac stimulant activity of bark and wood of *Premna integrifolia*. *Bangladesh J Pharmacol*. 2008; 3:107-113.
64. Rajendran R, Basha NS. Cardioprotective effect of ethanol extract of stem-bark and stem-wood of *Premna serratifolia* Linn. *Res J Pharm Tech*. 2008; 1:487-491.
65. Khatun H, Majumder R, Mamun A, Alam EK, Jami SI, Alam B. Preliminary pharmacological activity of the methanolic extract of *Premna integrifolia* barks in rats. *Avicenna J Phytomed*. 2014; 4:215-224.
66. Vadivu R, Jerad S, Girinath K, Kannan PB, Vimala R, Kumar NM. Evaluation of hepatoprotective and *in-vitro* cytotoxic activity of leaves of *Premna serratifolia* Linn. *J Sci Res*. 2009; 1:145-152.
67. Singh CR, Nelson R, Krishnan PM, Mahesh K. Hepatoprotective and anti-oxidant effect of root and root callus extract of *Premna serratifolia* L. in paracetamol induced liver damage in male albino rats. *Int J Pharma Bio sci*. 2011; 2:244-252.
68. Alamgir M, Rokeya B, Hannan JM, Choudhuri MS. The effect of *Premna integrifolia* Linn. on blood glucose in streptozotocin induced type 1 and type 2 diabetic rats. *Pharmazie*. 2001; 56:903-904.
69. Kar A, Choudhary BK, Bandyopadhyay NG. Comparative evaluation of hypoglycaemic activity of some Indian medicinal plants in alloxan diabetic rats. *J*

- Ethnopharmacol.* 2003; 84:105-108.
70. Gokani RH, Lahiri SK, Santani DD, Shah MB. Evaluation of immunomodulatory activity of *Clerodendrum phlomidis* and *Premna integrifolia* root. *Int J Pharmacol.* 2007; 3:352-356.
  71. Shukla V, Yadav D, Phulara SC, Gupta MM, Saikia SK, Pandey R. Longevity-promoting effects of 4-hydroxy-E-globularinin in *Caenorhabditis elegans*. *Free Rad Biol Med.* 2012; 53:1848-1856.
  72. Shukla V, Phulara SC, Yadav D, Tiwari S, Kaur S, Gupta MM, *et al.* Iridoid Compound 10-O-trans-p-Coumaroylcatalpol Extends Longevity and Reduces Alpha Synuclein Aggregation in *Caenorhabditis elegans*. *CNS Neurol Disord Drug Targets.* 2012; 11:1-9.
  73. Banu UK. Evaluation of neuroprotective effect in the experimental model of febrile seizure of the roots of *Premna serratifolia* Linn. International Herbal Conference. Bangalore, 2009. (Poster No. SNPS IHCB-09781).