



Evaluation of total polyphenols and antioxidant activity of African grape (*Lannea macrocarpa*) Fruits

Muhammad S^{1*}, Yahaya Lema S²

¹ Department of Chemistry, Sokoto State University, Nigeria

² Department of Biological Sciences, Sokoto State University, Nigeria

* Corresponding Author: **Muhammad S**

Article Info

ISSN (online): 2582-7138

Volume: 03

Issue: 05

September-October 2022

Received: 23-07-2022;

Accepted: 16-08-2022

Page No: 75-78

DOI:

<https://doi.org/10.54660/anfo.2022.3.5.3>

Abstract

The aim of the research work is to evaluate the polyphenols and antioxidant activity of African grape fruits. The methanolic extract was evaluated for its phenolic contents using Folin-Ciocalteu reagent, antioxidant activity using 1, 1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity, and flavonoid using sodium nitrite solution. The total phenolic content was 380 ± 2.11 mg/100g gallic acid equivalent (GAE), antioxidant activity of $562.25 \pm 2.91\%$, and flavonoids contents of 20.63 ± 1.80 mg/100g quercetin equivalent (QE). The results indicated the importance of *Lannea macrocarpa* fruits as potential source of polyphenols which if properly utilized could supplement the existing sources of antioxidants for the formulation of drugs and other therapeutic uses.

Keywords: *Lannea macrocarpa*, polyphenols, antioxidant, fruit pulp, flavonoids

Introduction

The global overpopulation needs parallel increase in food and nutrition sources. Food security becomes vulnerable when it is only dependent on a few numbers of traditional crop plants and domestic animals. Food and nutrition security need to be addressed in the context of biodiversity, an important asset to domesticate new crops or improve the quality of traditional crop plants (Hegazy *et al.*, 2013) [10]. Nutritionally, not only the quantity and energy contribution of foods are important to combat malnutrition but also their quality, including macro- and micronutrient content, and antioxidant activities. The gap between wild edible fruits and cultivated ones is wide and needs to be bridged by shedding more light on potential wild food biodiversity (FAO, 2010) [6]. Wild food plants represent a minor contribution to family meals, they are potentially important nutrient and cultural resources for local people around the world (Hegazy *et al.*, 2013) [10]. They often contain higher amount of nutrients and bioactive compounds than many cultivated species, especially those which have been under cultivation for many generations (Hegazy *et al.*, 2013) [10]. They have great potential as high-value nutraceuticals, and source of bioactive compounds for dietary supplements. Their fruits are edible and therefore important food items in traditional diets of local people, making an important contribution to the health of local communities. The edible fruits have been employed, for a long, in traditional and popular medicine (Delang, 2006) [4].

Fruits and vegetables are recommended as a source of dietary fiber, they are important part of a healthy diet, and variety is as important as quantity and no single fruit or vegetable provides all of the nutrients needed to be healthy. A diet rich in vegetables and fruits can lower blood pressure, reduce risk of heart disease and stroke, prevent some types of cancer, lower risk of eye defects, digestive problems and oxidative stress and also help the body to develop the capacity to fight against these by boosting immunity. This is based on the fact that they are home for many antioxidants such as ascorbic acid (vitamin C), tocopherols (vitamin E), carotenoids (provitamin A) and several phenolic compounds (flavones, isoflavones, flavanones, anthocyanins and catechins) (Shahidi and Nacz, 2004) [23].

African grape (*Lannea macrocarpa*) (Fig 1) commonly known as “Faaru” in Hausa speaking language belongs to the family *Anacardiaceae*. It is found in the savanna and the drier forest re-growth zone of West Africa. In Nigeria it is commonly found in Northern states of Sokoto, Kebbi, Zamfara, Kaduna, Katsina, Kano, and Jigawa. The unripe fruits are green in color while ripe ones are purplish black. The fruits are often sold in both city street markets and along roadsides in West Africa. In Nigeria, the tree is cultivated commercially on a small scale and the trees can be seen in and around villages. The fruit can be eaten fresh or dried like raisins for longer-term storage. The fruit makes an excellent jam, can be made into wine, and the pulp fermented into a potent alcoholic drink.



Fig 1: Ripped fruits of *Lannea macrocarpa*

Lannea macrocarpa is one of the most important species in Africa south of Sahara. The species belongs to the family of *Anacardiaceae*. It is a Soudano-Sahelian species also present in the Guineo-Congolese region. This species has a great socioeconomic importance in Burkina Faso (Abdoulaye *et al.*, 2014) [1]. It is a source of food for both human (fruits) and livestock (forage), and used in local handicrafts (woodcarving, dyeing, and tanning). The plant is a real panacea: the parts (leaves, bark and fruits) are used in the composition of diverse formula in traditional pharmacopoeia (Abdoulaye *et al.*, 2014) [1]. *Lannea macrocarpa* ripe fruit which is dark red is assumed to be an interesting food colorant. This color is due to the presence of flavonoids and anthocyanin pigments. Patricia *et al.* (2014) [19] reported that the fruits are eaten raw or dried, and a number of fermented or soft drinks are produced from its juice, however, important concentrations of phenolic compounds with strong antioxidant capacities have been found in the fruit (1005.75 mg/100 g of fruit). Partial reports on its seed and seed oil protein and lipid compositions are also available (Glew *et al.*, 1997).

Wild plant fruits contain many natural antioxidants compounds such as carotenoids, vitamins, phenols, tannins, flavonoids, and many secondary metabolites; which have been identified as a free radical or active oxygen scavengers (Oliveira *et al.*, 2009) [17]. Therefore, wild plant food sources need to be investigated not only from the nutritional value point of view, but also as a potential therapeutic agent against a wide range of human diseases (Oliveira *et al.*, 2009) [17].

Materials and Methods

Sampling and Sample treatment

Fresh fruits of African grape (*Lannea macrocarpa*) fruits were collected in September, 2021 from Chimola district of Gwadabawa Local government, Sokoto State, Nigeria. Five

(5) trees were randomly selected and only ripped fruits were collected from different branches of the trees as described by Hassan and Umar (2004) [9]. The sample was collected in black polythene bags and transported to the laboratory. Prior to analyses, the sample was authenticated at the Herbarium section, Botany Unit, Department of Biological Sciences, Usmanu Danfodiyo University, Sokoto, Nigeria. A representative sample was taken using alternate shovel method (Alan, 1996) [2]. The sample was thoroughly washed with distilled water and then air dried.

Preparation of the Extract

After drying, the fruits pulp was removed and crushed into powder with the help of pestle and mortar. Fifty grams (50g) of the powdered pulp were then soaked into 500cm³ methanol and allowed to stand for four days at 4⁰C. The extract was centrifuged at 1000rpm for 5minutes, filtered and then concentrated to dryness using rotary evaporator. The percentage extract was calculated using equation 1.

$$\% \text{ Extract} = \frac{\text{Weight of extract}}{\text{Sample weight}} \times 100 \dots \dots \dots (1)$$

The residue obtained was kept at 4⁰C until when required (Mothlanka *et al.*, 2012) [15].

Determination Total Polyphenols

The amount of total polyphenols in the sample was determined using modified Folin-Ciocalteu colorimetric method (Singleton *et al.*, 1999) [22]. Stock solution of sample extract (25 μ l) was dissolved in methanol and further dilution were performed to obtain readings within the standard curve made with gallic acid (R=0.997). The extract was oxidized by the Folin-Ciocalteu reagent (120 μ l) and the neutralization was made with Na₂CO₃ (340 μ l after 5 minutes). The absorbance was measured at 750 nm after 90 minute in the dark at room temperature. The result was expressed as milligram of gallic acid per 100 grams.

Determination of Flavonoids

The method reported by Kim *et al.* (2003) [13] was adopted. The methanolic extract (1.5mL) was added to 10mL volumetric flask filled with 5mL distilled water and 5% NaNO₂ followed by thorough mixing. To the content, 1.5mL of 2% methanolic AlCl₃ solution was added followed by 2mL of 1M NaOH solution and the volume made up to the mark with distilled water, the mixture was shaken vigorously and then incubated for 10minutes after which the absorbance measured at 367nm. The flavonoids content was calculated using a standard curve prepared from quercetin, and express as mg quercetin/100g of the extract.

Determination of Total Antioxidant Capacity

The total antioxidant capacity of the extract was determined by adopting the method reported by Pan *et al.* (2007). One mL of the extract was combined with 3mL reagent solution (0.6 M H₂SO₄, 28mM sodium phosphate and 4 mM ammonium molybdate). The reaction mixture was capped incubated in a thermal block at 95⁰C for 150 min after cooling at room temperature; the absorbance was measured at 695 nm against blank. Readings were taken every 30 min. The absorbance at 734 nm was measured to represent the total antioxidant activity and then calculated using equation 3.

$$\text{Total Antioxidant activity (\%)} = \left(1 - \frac{A_{\text{Sample}}}{A_{\text{Control}}}\right) \times 100 \dots \dots \dots (3)$$

Where A_{sample} and A_{control} represent the absorbance of the sample and control respectively.

Determination of DPPH Scavenging Activity

The reduction capacity of the DPPH radical was determined by the decrease of absorbance induced by antioxidants according to Hanane *et al.* (2017) [8]. The reaction system consisted of 0.1 mL of the extract and the standard diluted to different concentrations and 2.9 mL of a 0.025g/L DPPH in methanol. The mixture was shaken vigorously and left standing at room temperature in the dark for 30 minutes. The absorbance was measured at 515 nm against a blank. The ability to scavenge the DPPH radical was calculated using the formula in equation 4.

$$\% \text{ Scavenging effect} = \frac{(\text{ADPPH} - A_{\text{Sample}})}{\text{ADPPH}} \times 100 \dots \dots \dots (4)$$

Where A_{sample} is the absorbance of the solution when the sample extract was added while ADPPH represents the absorbance of the DPPH.

Statistical Analysis

The results obtained were statistically analyzed using one-way analysis of variance (ANOVA) with SPSS version 10.0 statistical packages and were reported as mean \pm standard error of mean.

Results

The result of the percentage yield, total polyphenols, flavonoids, antioxidant activity and DPPH scavenging activity of the fruits extract were expressed on dry weight basis (DW) and are presented in Table 1.

Table 1: Total polyphenols and Antioxidant activity of *Lannea macrocarpa* fruits

Parameter	Concentration
Yield	6.74 \pm 1.65 %
Total polyphenols	380 \pm 2.1 mgGAE/100g
Total flavonoids	20.63 \pm 1.80 mgQE/100g
Antioxidant activity	562.25 \pm 2.91 %
DPPH Scavenging activity	65.77 \pm 7.80 %

The values are mean \pm Standard error of mean; GAE = Garlic Acid Equivalent; QE = Quercetin Equivalent

Discussion

The Percentage yield

The percentage yield of the extract was 6.74 \pm 1.65g/100g of the fruit pulp which is an indication that the fruits contain some important nutritional and or medicinal phyto-compounds.

Total Polyphenols

The total polyphenols content of *Lannea macrocarpa* fruits pulp was 380 \pm 2.11 mgGAE/100g DW. The value recorded is lower than 424.84 \pm 20 mgGAE/100G DW for Straw berry, 398.25 \pm 0.1 mgGAE/100gDW for African star apple fruits, and higher compared to 247.25 \pm 11 mgGAE/100gDW for Black berry fruits (Ewa *et al.*, 2009) [5]. The value obtained is an indication that the fruits if properly utilized could be a

good source of polyphenols. Polyphenols are aromatics secondary plant metabolites that are widely spread throughout the plant kingdom and are associated with color, sensory qualities, nutritional and antioxidant properties (Robin, 2003). In food, polyphenols may contribute to the bitterness, astringency, color, flavor, odor and oxidative stability. Epidemiological studies and associated meta-analyses strongly suggested that long term consumption of diets rich in plant polyphenols offered some protection against development of cancer, cardiovascular diseases, diabetes, osteoporosis and neurodegenerative diseases (Kanti and Syed, 2009) [12].

Total Flavonoids

The flavonoids content of the fruits is 20.63 \pm 1.80 mgQE/100Gdw. The value is remarkably lower compared to 84.33 \pm 8mgQE/100Gdw for Straw berry and 29.07 \pm 1.12mgQE/100Gdw for Black berry (Andre *et al.*, 2011) [3] also lower than that of *Adansonia digitata* (42.73mgQE/100Gdw) reported by Lmien – Meda *et al.* (2008) [14]. The result obtained indicates that *Lannea macrocarpa* fruits are importance sources of flavonoids which are responsible for the attractive colors of flowers, fruits, leaves and also possess biological activities such as anti –inflammatory, anti-carcinogenic and anti-ant atherosclerotic activities (Olajire and Azeez, 2011) [16]. They are important in the plant for normal growth development and defense against infection and injury (Hasna El Gharas, 2009).

DPPH Scavenging and Antioxidant activities

2,2 – diphenyl -1 – picrylhydrazyl (DPPH) is a stable free radical that has been used as a tool to estimate the free radical scavenging activity of antioxidants. The *Lannea macrocarpa* fruits analyzed showed DPPH scavenging activity of 65.77 \pm 7.80 %. The value is higher compared to 46.64 \pm 1.65% reported for Blue berry fruits (Andre *et al.*, 2011) [3]. DPPH is one of the compounds that possessed a proton free radical with a characteristic absorption which decreases significantly on exposure to proton radical scavengers (Yamaguchi *et al.*, 1998). The DPPH scavenging by antioxidants is due to their hydrogen-donating ability. Antioxidants react with DPPH reducing a number of DPPH molecules equal to the number of available hydroxyl group (Hanane *et al.*, 2017) [8].

The antioxidant activity of the fruit extract (562.25 \pm 2.91 %) is probably due to its phenolic contents. It is a well – known fact that phenolic compounds are constituents of many plants, and have attracted great deal of public and scientific interest because of their health promoting effects as antioxidants (Hollman and Katan, 1999). The phenolic compounds exhibit considerable free radical scavenging activities through their reactivity as hydrogen or electron donating agents, and as metal ion chelating properties (Rice-Evans *et al.*, 1996) [20].

Conclusion

The study provides information on the phenolic composition, flavonoids content, and antioxidant activity. The results obtained indicated that the fruits if properly utilized can be a potential source of dietary polyphenols, and flavonoids which are important antioxidants and therefore their consumption should be stimulated based on the fact that the beneficial effects of polyphenols have been ascribed to their strong antioxidant activity and their ability to scavenge oxygen radicals and other reactive species. These features make

Lannea macrocarpa fruit a potentially interesting material for the development of functional foods or possible therapy for the prevention of some diseases.

Acknowledgments

The Author wishes to acknowledge the Tertiary Education Trust Fund (Tetfund) for funding the research through the Institution Based Research (IBR). The contributions of Professor S. M. Dangoggo of the Department of chemistry, Sokoto State University, Sokoto and Professor Sanusi Hassan Warra of the Department of Biochemistry, Usmanu Danfodiyo University, Sokoto is also highly appreciated for the success of this research work.

References

1. Abdoulaye Sereme, Jeanne Millogo, Sita Quinko, Mouhoussine Nacro. Micropropagation of a West Africa wild grape (*Lannea microcarpa*). International Journal of Biological and Chemical Sciences. 2014; 8(3):862-870.
2. Alan W. Soil and the environment: An introduction, Cambridge University press, 1996, 11-23.
3. Andre Bunea, Dmutrita O Rugina, Adele M Pinte, Zorita Sconta, Claudiu I Bunea, Carmen Socaciu. Comparative Polyphenolic Content and Antioxidant activities of some Wild and Cultivated Blueberries from Romania. Not. Bot. Horti Agrobi. 2011; 39(2):70-76.
4. Delang CO. Not just minor forest products: The economic rationale for the consumption of wild food plants by subsistence farmers. Ecology and Economy. 2006; 59:64-73.
5. Ewa Jabl Onska-Rys, Marta Zalewska-Korona, Janusz Kalbarczyk. Antioxidant Capacity, Ascorbic acid and Phenolic content in wild edible fruits. Journal of Fruit and Ornamental Plant Research. 2009; 17(2):115-120.
6. FAO/WHO. Trace elements in human nutrition and health (WHO/FAO/IAEA,) WHO, Geneva, 2010.
7. Glew RS, Vanderjagt DJ, Huang YS, Chuang LT. Nutritional analysis of the edible pit of *Sclerocarya birrea* in the Republic of Niger (daniya, Hausa). Journal of Food Composition and Analysis. 1995; 17:99-111.
8. Hanane B, Lilia B, Abdurrahmen R, Khaled A, Sami A, Mohammed K, et al. In vitro Cytotoxic and antioxidant activities of phenolic components of Algerian *Achillea odorata* leaves. Arabian Journal of Chemistry. 2017; 10:403-409.
9. Hassan LG, Umar KJ. Proximate and mineral composition of seeds and pulp of African locust bean (*Parkia biglobosa* L.) Nigerian Journal of Basic and Applied Sciences. 2004; 13:15-27.
10. Hegazy AK, Alruwaily SL, Faisal M, Alatar AA, El-Bana MI, Asaaeed AM. Nutritive value and antioxidant activity of some edible wild fruits in the Middle East. Journal of Medicinal Plant Research. 2013; 7(15):938-946.
11. Holiman PCH, Katan MB. Dietary flavonoids: Intake, health effects and availability. Food Chemistry and Toxicology. 1999; 37:937-942.
12. Kanti Bhooshan Pandey, Syed Ibrahim Rizvi. Plant polyphenols as dietary antioxidants in human health and disease. Oxidative Medicine and Cellular longevity. 2009; 2(5):270-278.
13. Kim DO, Jeong SW, Lee CY. Antioxidant capacity of phenolic phytochemicals from various cultivars of plums. Food Chemistry. 2003; 81:321-326.
14. Lamien-Meda ACE, Lamien MMY, Compaore RNT, Meda M, Kiendrebeogo B Zeba. Polyphenol content and antioxidant activity of fourteen wild edible fruits from Burkina Faso. Molecules. 2008; 13:581-594.
15. Motlhanka DM, SW Makhabu. Medicinal and edible wild fruit plants of Bostwana as emerging new crop opportunities, Journal of Medicinal Plants Research. 2012; 5(10):1836-1842.
16. Olajire AA, Azeez I. Total antioxidant activity, phenolic, Flavonoids, and Ascorbic acid contents of Nigerian vegetables. African Journal of Food Science and Technology. 2011; 2(2):22-29.
17. Oliveira I, Coelho V, Baltasar R, Pereira JA, Baptista P. Scavenging capacity of strawberry tree (*Arbutus unedo* L.) leaves on free radicals. Food Chemistry and Toxicology. 2009; 47:1507-1511.
18. Pan Y, Zhu J, Wang X, Zhang Y, He C, Ji X, et al. Antioxidant activity of some extract of Cortes fraxin and use in Peanut oil. Food Chemistry. 2007; 103:913-918.
19. Patricia Morales, Isabel CFR Ferreira, Ana Maria Carbalho, Virginia Fernandez-Ruiz M, Cortes Sanches, Montana Camara, et al. Wild edible fruits as a potential source of Phytochemicals with capacity to inhibit lipid peroxidation. European Journal of Lipid Science and Technology. 2014; 115:176-185.
20. Rice-Evans CA, Miller NJ, Paganga G. Structure antioxidant activity relationship of flavonoids and phenolic acids. Free radical Biology and Medicine. 1996; 20:933-956.
21. Robbins RJ. Phenolic acids in foods: an overview of analytical methodology. Journal of Agricultural Food Chemistry. 2003; 51:2866-2887.
22. Singleton VL, Rossi JA, Abel C. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. American Journal Enology Viticul. 1999; 16:44-158.
23. Shahidi F. Antioxidant properties of food phenolics. Phenolics in Food and Nutraceuticals. CRC Press, Boca Raton, FL, USA, 2004, 132-210.
24. Yagamuchi T, Takamura H, Matabo T, Terao J. HPLC method for evaluation of the free radical-scavenging activity of food by using 1, 1 di phenyl -2-picrylhydrazyl. Bioscience, Biotechnology, and Biochemistry. 1998; 62:1201-1204.