



Effects of salinity and basalt powder “Farina di Basalto®” interaction on seedlings of a white lupin variety

Yosr Ben Mhara ^{1*}, Anis Sakhraoui ², Yosr Ferchichi ³, Ichrak Kharbech ⁴, Giuliano Ragnoni ⁵, Gianluca Pizzuti ⁶, Fabio Primavera ⁷, Alessandro Riccini ⁸, Slim Rouz ⁹

^{1-4, 9} University of Carthage, Higher School of Agriculture of Mograne, Department of Agricultural Production, Laboratory of Agricultural Production Systems and Sustainable Development (SPADD), LR03AGR02, 1121 Mograne, Zaghuan, Tunisia

^{1, 3} University of Carthage, National Institute of Agronomy of Tunis, 43 Avenue Charles Nicolle, Tunis 1082, Tunisia

² University of Jendouba, Higher School of Agriculture of Kef, Le Kef 7119, Tunisia

⁵⁻⁸ Basalti Orvieto srl – Loc Cornale, 05014-CASTEL VISCARDO (TR), Italy

* Corresponding Author: Yosr Ben Mhara

Article Info

ISSN (online): 2582-7138

Volume: 03

Issue: 06

November-December 2022

Received: 04-11-2022;

Accepted: 25-11-2022

Page No: 508-512

Abstract

The alleviation of abiotic stresses such as salinity, is one of the most important challenges to respond to the increased food demand of the growing world population and to ensure sustainable agriculture.

This work aimed to investigate the interaction effects of sodium chloride (NaCl) and basalt powder “Farina di Basalto®” (FdB) on germination and early seedling growth parameters of *Lupinus albus* L. Combined treatments were evaluated, by applying four NaCl concentrations (0, 75, 150, 300 mM) on lupine seeds that were previously coated by five levels of “Farina di Basalto®” Type XF (0%, 1%, 3%, 5% and 10%).

Analysis of variance showed that salinity had a significant negative effect on germination percentage and the studied parameters during the juvenile stage ($p < 0.05$). This effect depended on the different NaCl levels. The treatment with “Farina di Basalto®” Type XF showed a positive effect on growth parameters. Results revealed that in comparison with control, different concentrations of “Farina di Basalto®” Type XF stimulated hypocotyl elongation to reach the higher increase by 131% compared to control, under 150 mM NaCl with 10% of “Farina di Basalto®” Type XF. In addition, Basalt in combination with salt stress have a significant effect on the salinity tolerance index and revealed the best results in salinity resistance in the presence of 10% of “Farina di Basalto®” Type XF.

Keywords: Salt stress, “Farina di Basalto®”, germination, early seedling, *Lupinus albus* L

Introduction

White lupin (*Lupinus albus* L.), belongs to the Fabaceae family. It is cultivated for human consumption, green manuring and as forage. This grain legume has Mediterranean origins, where drought, salinity and mineral deficiency are among the major constraints for lupines production ^[1]. Hence the interest of evaluating salt stress tolerance of white lupine to develop this culture in different regions threatened by soil salinization.

An increase in the salinity concentration in the soil solution leads to an increase in osmotic pressure, which necessitates an increased effort of the plants to absorb the nutrients, which negatively affected their growth and production ^[2, 3].

Plants are able to detoxify excessive oxidation caused by salt-stress conditions by complex enzymatic and nonenzymatic mechanisms that protect plant cells against oxidative damage ^[4].

The basalt is a tuff originates from volcanic projections ^[5]. It is sometimes observed as blocks or even ashes with black, red or even dark green color. This volcanic igneous rock is generally employed in different areas such as construction, industrial engineering and agronomy ^[6, 7]. In fact, the basalt is known as a natural fertilizer used to improve soil chemical proprieties ^[8]. This rock contains many chemical components required in agronomy and soil fertilization such as SiO₂ (37.76 to 59.64%), Al₂O₃ (11.77 to 14.32%), CaO (5.57 to 14.75%), MgO (5.37 to 9.15%), Fe₂O₃ (10.1 to 20.93%), K₂O (1.7 to 6.69%), Na₂O (1.4

to 3.34%) and TiO₂ (1.81 to 3.73%)^[8]. This mineral used to restore soils fertility, has shown also a positive effect on plant growth, total yield and fruits quality^[9,10]. Besides, basalt was shown in many studies to confer resistance against pests and diseases to several crops, by stimulating their natural defense reactions in fields and even under storage conditions^[8, 11,12]. Furthermore, an alternative strategy to decrease the oxidative damage on plants is to use silicon (Si) fertilization, the principal component of Basalt powder, in plants subjected to environmental stresses^[13]. Therefore, Si has been shown to be able to improve early growth and establishment of plants under stressful conditions through increasing antioxidant enzymes, photosynthetic capacity and low transpiration coefficient^[14, 15]. Which incites studying the effect of Basalt powder on improving cultivated species tolerance to salt stress.

This study aims to evaluate the capacity of “Farina di Basalto®” to alleviate salinity effects on a white lupine variety, at the germination phase and early seedling growth by measuring various parameters under controlled conditions.

Material and methods

Plant material

Seeds of *L. albus* were obtained from the lupine collection of the Genetic Diversity and Molecular Characterization Laboratory of Higher School of Agriculture of Mograne.

Basalt powder “Farina di Basalto®”

Basalt powder or “Farina di Basalto®” type XF, produced by Basalti Orvieto s.r.l, was used in this work. It's composed of micronized particles with less than 20 µm obtained through an industrial process. The raw material used for the production of “Farina di Basalto®” comes from the Castel Viscardo deposit (TR) - Italy. It is a basic volcanic rock, with a vitreous structure, which does not contain free crystalline silica, nor amentiferous minerals or other substances harmful to the environment or animal health. The micronized “Farina di Basalto®” is obtained by the mechanical grinding of the pure mineral, using ceramic tools, without the addition of other minerals or substances, nor the use of materials containing washing water with flocculants, or undesirable and harmful products in agriculture. Known for its fertilizing effects applied in agriculture, “Farina di Basalto®” contains natural elements such as Silicon, Alumina, Potassium and Calcium^[16]. Silica (silicon dioxide) is the main component that characterizes Basalt Flour(powder) with percentages ranging from 45% to 49%^[12]. Physicochemical characteristics and the different components of Basalt powder “Farina di Basalto®” type XF” are presented in the table 1 and table 2 respectively.

Table 1: Physicochemical characteristics of Basalt powder “Farina di Basalto®” type XF”^[12].

Physicochemical characteristics	
Solid physical state	Powdery
Water solubility	Not soluble in water
Color	Slightly gray
Odor	Not noticeable
pH	9±0.5 (Log[H+])
Electrical conductivity (2:1 extract)	1.14 (dS /m)
Cation exchange capacity	9 (meq/100g)
Assimilable iron (As. Fe)	377 (mg/kg)
Density	2.70 (kg/dm ³)

Table 2: Chemical proprieties of Basalt powder “Farina di Basalto®” type XF”^[12].

Chemical composition (%)	
SiO ₂	45-49
Al ₂ O ₃	20.5-25.6
K ₂ O	8-10
Fe ₂ O ₃	5.2-8.5
CaO	7.5-8.5
MgO	1.9-2.6
Na ₂ O	2.2-4.9
P2O5	0.6-0.7

Experimental design

This study was carried out in the Genetic Diversity and Molecular Characterization Laboratory of Higher School of Agriculture of Mograne located at 156 m of altitude, 10.092049° of longitude and 36.428272° of latitude. Randomly selected seeds were disinfected using 5% sodium hypochlorite for 3 min, then thoroughly rinsed with distilled water to remove sodium hypochlorite and placed in sterilized 9 cm ø petri dishes upholstered by two layers of Whatman® n°1 filter paper (5 seeds per petri dishes). Germination and juvenile growth were rated under four concentrations of sodium chloride and five basalt powder percentage, with four replicates for each treatment. Basalt powder, lightly moistened with distilled water, was mixed with disinfected seeds using five percentages 0, 1, 3, 5 and 10% corresponding to 0, 3.57, 10.73, 17.88 and 35.77 g/Kg according to the weight of the sampled seeds. Salt stress was applied by subjecting lupine seeds previously coated with different parentages of basalt powder, to 10 ml salt solutions of 75, 150 and 300 mM NaCl. Distilled water was used as control. Petri dishes were placed in the dark in an incubator at 20±1 °C and 40 % relative humidity for ten days.

Studied parameters

Seed germination was recorded every 24 h for 10 days. A seed was considered germinated when the radicle reaches 2 mm length^[17]. At the end of the experiment, final germination percentage (GP) was calculated according to the formula cited by El Rasafi^[18].

$$GP = \frac{\text{Germinated seeds}}{\text{Total seeds}} \times 100$$

Hypocotyl and radicle length were measured after 10 days, using a digital calliper. Seedlings fresh and dry weight were subsequently recorded. The dry weights were measured after drying fresh seedlings in a laboratory oven (KOTERMANN 2771, Uetze, Germany) at 70°C for 48h.

Salt tolerance index STI was calculated following the formula of Aghamir *et al*^[19].

$$STI = \frac{\text{Seedling dry weight of stressed plant}}{\text{Seedling dry weight of control plants}}$$

Statistical analysis

A two-way analysis of variance (ANOVA) was carried out to test the combined effect of four salt stress treatments (0, 75, 150, 300 mM) and five basalt powder concentrations (0, 1, 3, 5, 10 %) on germination and early seedling growth of *L. albus*. Variation of data was statistically significant for Tukey's multiple range tests at the $p < 0.05$ level.

The statistical analysis of this data was done using 'PLAnt Breeding STATistical software' (PLABSTAT), version 3A of 2011-06-14 [20].

Results and discussion

Results presented in table 2 showed a significant effect of salinity and basalt treatments on the germination percentage of *Lupinus albus* seeds ($p < 0.05$). In fact, a decrease of germination percentage was observed proportionally to the increase of NaCl concentrations as well as for the lowest basalt levels 1% and 3%. While, for basalt treatment, a significant re-increase of germination percentage was recorded at the rate of 5%, and this for all studied salinity concentration with an average of 83.75%. The interaction between the two treatments was non-significant for this parameter. Shoot length and radicle length decreased along with increasing salinity (Figure1). "Farina di Basalto" type XF showed a positive effect on these growth parameters for control seeds, with an increase of shoot growth by 12% at 1% basalt level.

However, under salt stress, this increase of 12% is reached at 10% basalt treatment in the presence of a low concentration of sodium chloride 75 mM. Analysis of variance revealed a significant interaction ($p < 0.05$) between salinity and basalt treatments on cotyledon length of *L. albus* seedlings (Table 2). A significant enhancement of cotyledon length of stressed seedlings, was demonstrated, in presence of 10% of "Farina di Basalto" type XF". At 300 mM NaCl, cotyledon length increased by about 5.5 % at 10% of basalt treatment compared to control. On the other side, FdB application stimulated hypocotyl and radicle length proportionally to the increasing basalt levels for all sodium chloride concentrations. Results recorded that hypocotyl and radicle lengths average increased by 49.8 % and 17% respectively, at 10 % of basalt treatment when compared to control. The influence of 10% of FdB was more effective on hypocotyl length at 75 mM and 150 mM NaCl, by an increase of 55 % and 131 % respectively compared to control.

Table 2: Combined effect of salinity and "Farina di Basalto" Type XF" treatments on germination and seedling characteristics of *Lupinus albus*.

Salinity (mM) **	"Farina di Basalto® Type XF" treatment (%) **					
	0	1	3	5	10	Means
	Germination percentage (%)					
0	100	100	90	95	85	94.00 ^a
75	95	90	90	95	95	94.00 ^a
150	90	85	75	90	70	84.00 ^a
300	80	53.3	35	50	15	47.67 ^b
Means	93.75 ^a	82.08 ^{ab}	72.50 ^b	83.75 ^{ab}	67.50 ^c	79.92
	Cotyledon length (mm)					
0	24.51±0.23	25.04±0.66	18.26±1.26	21.5±2.61	23.41±0.28	22.54 ^{a±1}
75	23.58±0.44	19±1.47	21.34±2.2	20.03±0.97	23.69±0.28	21.53 ^{ab±1.07}
150	22.83±0.52	19.69±2.23	15.08±2.10	19.04±2.35	23.5±0.35	20.10 ^{b±1.08}
300	20.19±1.66	10.31±2.06	7.1±2.87	22.23±0.55	21.31±0.66	16.23 ^{c±1.56}
Means	22.78 ^{a±0.46}	18.5 ^{b±1.60}	15.45 ^{c±2.10}	20.79 ^{ab±6.48}	22.98 ^{a±0.39}	20.10±1.69**
	Hypocotyl length (mm)					
0	34.53±2.79	40.92±2.54	27.03±2.21	36.22±6.11	41.16±2.52	35.97 ^{a±3.17}
75	23.52±3.58	29.51±4.11	25.07±3.02	32.51±3.39	36.64±2.27	29.45 ^{a±3.27}
150	11.29±1.64	19.11±5.20	18.62±2.96	21.14±2.21	26.1±1.92	19.25 ^{b±2.79}
300	0	0	0	0	0	0
Means	23.11 ^{c±2}	29.85 ^{b±2.96}	23.57 ^{c±2.04}	29.96 ^{b±2.92}	34.63 ^{a±1.14}	28.22±2.03
	"Farina di Basalto® Type XF" treatment *					
	Fresh weight (mg)					
0	3.31±0.047	3.77±0.14	2.62±0.14	2.95±0.33	3.34±0.15	3.20 ^{ab±0.16}
75	2.82±0.11	4.76±1.81	2.91±0.27	2.98±0.15	3.7±0.06	3.43 ^{a±0.48}
150	2.7±0.08	2.72±0.35	2.14±0.33	2.66±0.42	3.11±0.13	2.67 ^{b±0.26}
300	2.41±0.23	1.25±0.25	0.97±0.27	2.62±0.14	2.61±0.18	1.97 ^{c±0.21}
Means	2.81 ^{a±0.11}	3.12 ^{a±0.63}	2.16 ^{b±0.25}	2.80 ^{ab±0.26}	3.19 ^{a±0.13}	2.82±0.27
	"Farina di Basalto® Type XF" treatment					
	Dry weight (mg)					
0	0.74±0.05	0.79±0.04	0.78±0.02	0.81±0.03	0.77±0.02	0.78 ^{c±0.03}
75	0.79±0.02	0.81±0.04	0.83±0.02	0.85±0.02	0.92±0.02	0.84 ^{b±0.02}
150	0.88±0.01	0.9±0.04	0.92±0.02	0.92±0.01	0.91±0.03	0.91 ^{a±0.02}
300	1.03±0.12	0.95±0.03	0.92±0.03	0.78±0.07	0.91±0.04	0.9 ^{a±0.05}
Means	0.86 ^{a±0.05}	0.86 ^{a±0.02}	0.86 ^{a±0.2}	0.84 ^{a±0.03}	0.86 ^{a±0.02}	0.86 ^{*±0.02}
	Salt tolerance index					
0	1.00±0.07	1.07±0.034	1.06±0.03	1.10±0.05	1.04±0.16	1.06 ^{a±0.06}
75	1.07±0.05	1.09±0.02	1.12±0.03	1.16±0.03	1.24±0.04	1.14 ^{ab±0.03}
150	0.88±0.03	0.9±0.05	0.92±0.03	0.92±0.02	0.91±0.05	1.23 ^{b±0.03}
300	1.39±0.04	1.29±0.03	1.25±0.02	1.05±0.04	1.18±0.09	1.23 ^{b±0.04}
Means	1.16 ^{a±0.04}	1.17 ^{a±0.03}	1.17 ^{a±0.02}	1.14 ^{a±0.03}	1.17 ^{a±0.08}	1.16 ^{*±0.04}

Each value represents the mean ± SE (n = 4). Results of significant differences between accessions and treatments determined by two-way ANOVA (Tukey test, $p < 0.05$) are indicated by: *: $p < 0.05$; **: $p < 0.01$




Figure 1: Combined effect of salinity and “Farina di Basalto® Type XF” treatments on shoot and radicle lengths of *L. albus*. Bars represent mean \pm S.E. (n = 4). Different letters indicate a significant difference of treatments ($p < 0.05$, Tukey’s multiple range test)

L. albus seedlings fresh and dry weights vary according to the applied salinity levels. Fresh weight average recorded a significant decrease by 38.5 % at 300mM NaCl compared to control. However, dry weight average showed a significant increase by 16.6% for 300 mM of salt concentration when compared to control. Although the combined salinity and basalt treatments had a significant effect on dry weight, the different levels of basalt do not show a statistically significant difference on this parameter.

Results showed a significant increase in salt tolerance index when 10 % of basalt powder is applied to seedlings exposed to 75 mM of sodium chloride by 15%, nevertheless STI drops by 15% under 300 mM NaCl, when compared to control. These results confirm that coating seeds with 10 % of basalt powder alleviate especially low levels of salinity stress in white lupine at early growth phase.

These findings agree with previous studies. In fact, the increase of salt concentration induces a high osmotic pressure, thus reducing the amount of available water to be absorbed by the seeds or by the presence of toxic ions in the solution, which negatively affects germination process and seedling growth [21, 22].

Regarding Basalt, its addition had a positive impact on growth parameter. Since the main component of basalt is silicon, this result is in line with previous reports, which have suggested that silicon has many positive effects on the growth and yield as well as physiology and metabolism in different plant species [23, 24, 25]. In addition, the study conducted by Zuccarini [26] to evaluate the effect of silicon on *Phaseolus vulgaris* L. under salt stress showed that Si application enhanced stressed plants growth significantly and that salinity alleviation was more effective at lower NaCl (30mM) concentration. Those positive results were observed in our study at 75 mM and 150 mM, especially by the stimulation of hypocotyl elongation according to the increase in “Farina di Basalto®” percentage, which demonstrates a high tolerance of this white lupine variety to salinity and the effectiveness of the studied basalt levels with the best results registered at

10%. Other studies have proven the beneficial effects of silicon on increasing leaf area and photosynthesis by decreasing the decomposition of photosynthetic pigments as well as antioxidant enzymatic activity under stress conditions [25, 27, 28, 29].

Conclusion

Seed treatment with Basalt Powder, rich in silicon, may increase significantly salt stress tolerance in white lupine at the juvenile growth stage. The present research highlighted a new method to alleviate salinity stress using a natural rock, which constitutes an ecological solution to increase white lupine production and allow to this crop to cope with abiotic constraints. The effectiveness of “Farina di Basalto®” for other lupine species as well as the optimal doses to use under controlled conditions and in saline soil may be the subject of further studies.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgement

This work was performed at the Laboratory of Agricultural Production Systems and Sustainable Development. Authors wish to thank the head of SPADD laboratory, Professor Lamia Ajili Ghezal, and all the laboratory staff.

References

- Hussien ET. Salinity stress affecting viability and genetic stability of *Lupinus albus* L. *Vegetos*. 2022; 35:674-680.
- Parlak AÖ. Effect of salinity in irrigation water on some plant development parameters of sainfoin (*Onobrychis viciifolia* Scop.) and soil salinity. *Journal of Agricultural Sciences*. 2008; 14(4): 320-325.
- Mostafavi K, et al. Effect of salt stress on germination and early seedling growth stage of sugar beet cultivars. *American-Eurasian Journal of Sustainable Agriculture*. 2012; 6(2):120-125.

4. Alves RC, Medeiros A S, Nicolau MCM, Pizolato Neto A, Oliveira F A, Lima LW, *et al.* The partial root-zone saline irrigation system and antioxidant responses in tomato plants. *Plant Physiology and Biochemistry*. 2018; 127:366-379.
5. Rahmouni H, Ouari MT. Caractérisation et étude des propriétés d'usage des tufs de Remila. Mémoire fin d'étude. Université Abderrahmane Mira de Bejaia, Bejaia, 2016, 31.
6. Drobot NF, Noskova OA, Steblevskii AV, Fomichev SV, Krenev KA. Use of chemical and metallurgical methods for processing of gabbro-basalt raw material. *Theoretical Foundations of Chemical Engineering*. 2013; 47(4):484-8.
7. Pisciotta A, Perevozchikov BV, Osovetsky BM, Menshikova EA, Kazymov K P. Quality Assessment of Melanocratic Basalt for Mineral Fiber Product, Southern Urals, Russia. *Natural Resources Research*. 2014; 24:329-337.
8. Elimem M, Kalboussi M, Lahfeg C, Rouz S, Kharroubi H, *et al.* Evaluation of insecticidal efficiency of Basalt powder "Farina di Basalto®" to control *Tribolium castaneum* (Coleoptera; Tenebrionidae), *Rhyzopertha dominica* (Coleoptera; Bostrichidae) and *Ephestia kuehniella* (Lepidoptera; Pyralidae) on stored wheat. *IOSR Journal of Agriculture and Veterinary Science*. 2021; 14(5):01-06.
9. Fawzy ZF, El-Bassiony AM, Yunsheng L, Zhu O, Ghoname AA. Effect of mineral, organic and bio-N fertilizers on growth, yield and fruit quality of sweet pepper. *Journal of Applied Sciences Research*. 2012; 8(8):3921-3933.
10. Rouz S, Elimem M, Kharroubi H, Mekni A, Kouki S, *et al.* Effect of basalt powder « Farina di Basalto® » on pepper crop growth parameters under greenhouse. *IOSR Journal of Agriculture and Veterinary Science*. 2020; 13(11):29-39.
11. Isnugroho K, Hendronursito Y, Birawidha DC. Characterization and utilization potential of basalt rock from East-Lampung district. *Mineral Processing and Technology International Conference*. 2017; 285:1-5.
12. Anonymous. Scheda Tecnica dei principali elementi Caratteristiche fisico chimiche e composizione chimica Farina Di Basalto®. 2022.
13. Alves RC, Nicolau MCM, Checchio MV, *et al.* Salt stress alleviation by seed priming with silicon in lettuce seedlings: an approach based on enhancing antioxidant responses. *Bragantia*. 2020; 79:19-29.
14. Sivanesan I, Son MS, Lim CS, Jeong BR. Effect of soaking of seeds in potassium silicate and uniconazole on germination and seedling growth of tomato cultivars, Seogeon and Seokwang. *African Journal of Biotechnology*. 2011; 10:6743-6749.
15. Imtiaz M, Rizwan MS, Mushtaq MA, Ashraf M, Shahzad SM, Yousaf B, *et al.* Silicon occurrence, uptake, transport and mechanisms of heavy metals, minerals and salinity enhanced tolerance in plants with future prospects: A review. *Journal of Environmental Management*. 2016; 183:521-529.
16. Elimem M, Jaouadi R, Kalboussi M, *et al.* Management of *Ceratitis Capitata* and *Phyllocnistis citrella* with Basalt powder "Farina di Basalto®" compared to two botanical extracts (Citrus aurantium and Nerium oleander) in citrus orchard. *International Journal of Zoological and Entomological Letters*. 2022; 2(1):71-80.
17. Sayar R, Bchini H, Mosbahi M, *et al.* Effects of salt and drought stresses on germination, emergence and seedling growth of durum wheat (*Triticum durum* Desf.). *Journal of Agricultural Research*. 2010; 5(15):2008-2016.
18. El Rasafi T, Nouri M, Bouda S, Haddioui A. The Effect of Cd, Zn and Fe on Seed Germination and Early Seedling Growth of Wheat and Bean. *Ekológia (Bratislava)*. 2016; 35(3):213-223.
19. Aghamir F, Bahrami H, Malakouti MJ, Eshghi S, Sharifi F. Seed germination and seedling growth of bean (*Phaseolus vulgaris*) as influenced by magnetized saline water. *Eurasian Journal of Soil Science*. 2016; 5(1): 39-46.
20. Utz, H.F. Plabstat: A Computer Program for Statistical Analysis of Plant Breeding experiments. Universität Hohenheim; Stuttgart, Germany. Version 3A, 2011.
21. Pearson KE, Bauder JW. The basics of salinity and sodicity effects on soil physical properties. *Water quality and irrigation management*, 2003, 1-9.
22. Othman Y, Al-KaraKi G, Al-Tawaha AR, AlHorani A. Variation germination and ion uptake in genotype barley under salinity condition. *World Journal of Agricultural Sciences*. 2006; 2(1):11-15
23. Liang YC, Sun W, Zhu YG, Christie P. Mechanisms of silicon mediated alleviation of abiotic stress in higher plants: a review. *Environmental Pollution*. 2007; 147:422-428.
24. Ma JF, Yamaji N. Functions and transport of silicon in plants. *Cellular and Molecular Life Sciences*. 2008; 65:3049-3057.
25. Bybordi A. Effect of ascorbic acid and silicium on photosynthesis, antioxidant enzyme activity, and fatty acid contents in canola exposure to salt stress. *Journal of Integrative Agriculture*. 2012; 11(10):1610-1620.
26. Zuccarini P. Effects of silicon on photosynthesis, water relations and nutrient uptake of *Phaseolus vulgaris* under NaCl stress. *Biol. Plantarum*. 2008; 52:157-160.
27. Liang YC, Chen Q, Liu Q, Zhang W, Ding R. Exogenous silicon (Si) increases antioxidant enzyme activity and reduces lipid peroxidation in roots of salt-stressed barley (*Hordeum vulgare* L.). *Journal of plant physiology*. 2003; 160(10):1157-1164.
28. Al-aghaby K, Zhu Z, Qinhu S. Influence of silicon supply on chlorophyll content, chlorophyll fluorescence and antioxidative enzyme activities in tomato plants under salt stress. *Journal of Plant Nutrition*. 2005; 27:2101-2115.
29. Gong H, Zhu X, Chen K, Wang S, Zhang S. Silicon alleviates oxidative damage of wheat plants in pots under drought. *Plant Science*. 2005; 169:313-321.