



International Journal of Multidisciplinary Research and Growth Evaluation.

What to look for when trying to detect a viral pathogen?

Carlos Navarro Venegas

Associate Professor according to FAVET, Faculty of Veterinary and Livestock Sciences (FAVET), University of Chile, Santiago, Chile

* Corresponding Author: **Carlos Navarro Venegas**

Article Info

ISSN (online): 2582-7138

Volume: 04

Issue: 02

March-April 2023

Received: 26-01-2023;

Accepted: 16-02-2023

Page No: 25-26

Abstract

Molecular virology has been a multifaceted activity for some years backed by a professional practice of almost 30 years. In it we have incorporated several molecular biotools in the study of disease-causing pathogens in the area of human and animal medicine. According to André Lwoff: Viruses are viruses, so their detection does not vary much between the two medicines, as we will see below.

Keywords: viral detection, biotools, primer design, softwares

Introduction

There is no doubt that the fantastic idea of Kary Mullis and other scientists has allowed molecular virology, in addition to being a dynamic discipline, to have incorporated novel methodologies currently in use.

Viruses have special genomes (RNA or DNA) in terms of size, length, or nucleotide sequence. We can easily find these sequences in a special database: the Genbank®^[1], which together with some computer programs allow both the design of primers for a PCR reaction and the corroboration of the identity of the pathogen of interest.

Thus, today it is not possible to have any pretext for not trying to detect a pathogen of interest, whether it affects humans or another animal species and also to propose and give away some methodologies using computer programs even free and online. In a third world country like we could, so you too...!!!!!!

Material and methods

The brilliant idea of Kary Mullis^[2], called Polymerase Chain Reaction in concomitance with 2% agarose gel electrophoresis allows to observe the amplification of a DNA fragment of an expected size (conventional PCR). If this fragment is subsequently sent for sequencing and we use some computer programs such as CLUSTAL Omega^[3] and BLAST^[4], the nucleotide identity of the suspected agent is determined.

When considering the existing information in the Genbank® together with the program Oligoperfect from Invitrogen® or another similar one, which generates the best primers for the detection of any pathogen by PCR, confirms A.Lwoff's statement and that there is also no pretext for proposing a viral detection protocol, for example.

Another thing is with a guitar, they say, but having samples to carry out the PCR will never be the problem, because from fragments of organs or fluids from which the nucleic acid is extracted. Now, once again, thanks to Clustal Omega and BLAST it is possible to establish the identity of the pathogen, or at least a certain approximation.

Discussion

Undoubtedly, several of the methodologies correspond to scientific innovations and brilliant ideas such as that of Kary Mullis that together with biotools such as Clustal,

BLAST and other similar ones that have been mentioned in various publications in mainstream journals^[5,6] and others of equal importance that have magnified the molecular study of pathogens of veterinary interest^[7-17].

It is not for nothing that the brilliant idea of PCR is considered a level of the discovery of the structure of DNA or the theory of relativity, within the scientific legacies in the history of humanity.

Conclusion

Not everything is science for science's sake. Yes, our country has -at least- 35 new medical veterinarians who started without knowing these techniques and today could perfectly establish and develop a molecular methodology for the detection and diagnosis of a viral pathogen.

Acknowledgments

We thank the students of FAVET for making their dreams come true with us and Dr. Aron Mosnaim, from the Wolf Foundation, Illinois, USA (since 2020).

References

1. Genbank®. NIH genetic sequence database. 2023. Available from: <https://www.ncbi.nlm.nih.gov/genbank/>
2. Mullis K, Faloona F. Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. *Methods in Enzymology*. 1987; 155:335-350.
3. Clustal Omega. Multiple Sequence Alignment, 2023. Available from: <https://www.ebi.ac.uk/Tools/msa/clustalo/>.
4. Blast. Basic Local Alignment Search Tool, 2023. Available from: <https://blast.ncbi.nlm.nih.gov/Blast.cgi>
5. Vergara-Wilson V, Hidalgo-Hermoso H, Sánchez C, Abarca M, Navarro C, Celis-Diez S, *et al.* Canine Distemper Outbreak by Natural Infection in a Group of Vaccinated Maned Wolves in Captivity. *Pathogens*, 2021. Available from: <https://doi.org/10.3390/pathogens10010051>.
6. Hidalgo-Hermoso E, Cabello J, Vega C, Kroegeer-Gómez H, Moreira-Arce D, Napolitano C, *et al.* An eight-year survey for canine distemper virus indicates lack of exposure in the endangered Darwin's fox (*Lycalopex fulvipes*). *J Wild Dis*, 2020. Available from: <https://pubmed.ncbi.nlm.nih.gov/31833816/>.
7. Jara P, Céspedes P, Navarro C. Canine Distemper Virus detection based in Hemagglutinine gen as target in reverse transcriptase-Polymerase Chain reaction. *IVS*, 2018. Available from: <https://www.heighpubs.org/hvsr/pdf/ivs-aid1012.pdf>.
8. Salas V, Pizarro J, Navarro C. Phylogenetic analysis of canine distemper virus detected in Chile. *IJCR*. 2018; 10:72402-72407.
9. Gallegos M, Céspedes P, Pizarro J, Navarro C. Is the M gene of Canine Distemper virus a eligible target for detection? *EASJALS*, 2018. Available from: https://www.easpublisher.com/media/articles/EASJALS_11_22-32_c.pdf (P18).
10. Méndez-Valenzuela VK, Jara MA, Navarro C. Canine Distemper Virus: Multiple detection of the H and N genes by the Polymerase Chain Reaction associated with Reverse Transcriptase. *CJVDS*, 2021. Available from: <https://www.corpuspublishers.com/assets/articles/article-pdf-158.pdf>.
11. Vera C, Jara MA, Navarro C. A preliminary studio for the F gen of Canine Distemper Virus as target for phylogenetic analysis. *GSCARR*, 2022. Available from: <https://gsconlinepress.com/journals/gscarr/sites/default/files/GSCARR-2022-0034.pdf>.
12. Correa V, Céspedes PF, Navarro C. Promising use of Polymerase Chain Reaction Associated to Reverse Transcription for the Detection of the America-1 Lineage of Canine Distemper Virus. *IJRSZ*, 2019. Available from: <https://www.arcjournals.org/pdfs/ijrsz/v5-i1/4.pdf>.
13. Navarro C, Muñoz C, Céspedes P. The Nucleocapside Protein Gene As Excellent Target For Detection Of Canine Distemper Virus by Reverse Transcriptase-Polymerase Chain Reaction. *Am J Biomed Sci & Res.*, 2019. Available from: <https://biomedgrid.com/pdf/AJBSR.MS.ID.000935.pdf>
14. Mateo F, Céspedes PF, Navarro C. The Phosphoprotein Gene from Canine Distemper Virus as Target in Viral Detection. *IJZAB*, 2019. Available from: <https://medwinpublishers.com/IJZAB/the-phosphoprotein-gene-from-canine-distemper-virus-as-target-in-viral-detection.pdf>.
15. Bolívar P, Céspedes P, Navarro C. Use of the Reverse Transcription-Polymerase Chain reaction for differential detection of two lineages of the canine distemper virus in Chile. *IVS*, 2019. Available from: <https://www.heighpubs.org/hvsr/pdf/ivs-aid1014.pdf>.
16. Abarca MJ, Hidalgo E, Raggi LA, Navarro C. Genotypic evidence of infection by Canine Distemper Virus in maned Wolf from a zoological collection. *IJSR*, 2018. Available from: <https://www.ijser.org/researchpaper/Genotypic-evidence-of-infection-by-Canine-Distemper-Virus-in-maned-wolf-from-a-zoological-collection-in-Chile.pdf>
17. Pincheira D, Céspedes P, Pizarro J, Navarro C. Molecular detection of Canine Distemper Virus through the use of the Large Polymerase Gene. *EASJALS*, 2018. Available from: https://www.easpublisher.com/media/articles/EASJALS_12_39-43_c.pdf.