



## Community structure of mangrove ecosystem in Dulangeya Village, Gorontalo province

Rizkiyah Magfira Lasoma <sup>1\*</sup>, Pujiono Wahyu Purnomo <sup>2</sup>, Suryanti <sup>3</sup>

<sup>1-3</sup> Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Diponegoro University, Indonesia

\* Corresponding Author: **Rizkiyah Magfira Lasoma**

---

### Article Info

**ISSN (online):** 2582-7138

**Impact Factor:** 5.307 (SJIF)

**Volume:** 04

**Issue:** 06

**November-December 2023**

**Received:** 10-09-2023;

**Accepted:** 12-10-2023

**Page No:** 408-412

### Abstract

Dulangeya Village is one of the coastal villages in Boalemo Regency, Gorontalo Province which has mangrove ecosystem potential. This study aims to identify mangrove species and analyze the community structure of mangrove ecosystems in Dulangeya Village, Botumoito District, Boalemo Regency, Gorontalo Province. This research was conducted in January-March 2023. Data taken include species density, relative density, species frequency, relative frequency, species closure, relative closure, and importance value index. The method used was direct observation. Data collection on mangrove vegetation was carried out using 100m long line transects at 4 stations. Each transect consists of plots with a size of 10x10m. The results showed that there were 7 types of mangroves found in Dulangeya Village. Tree level density is classified as moderate and for sapling and seedling level is classified as good. The highest species frequency for tree level is *Rhizophora apiculata*, while for sapling and seedling level is *Ceriops decandra*. The Importance Value Index (IVI) for the tree level obtained values ranging from 5.247 - 127.159. This shows that mangrove trees have a fairly important value for the lives of the people of Dulangeya Village.

**DOI:** <https://doi.org/10.54660/IJMRGE.2023.4.6.408-412>

**Keywords:** Dulangeya Village, Important Value Indeks, Mangrove

---

### Introduction

Mangrove ecosystems are one of the marine resources that have many benefits and functions for the environment and humans (Karimah, 2017) <sup>[7]</sup>. Mangrove ecosystems have a role in controlling the environmental quality of coastal areas, including functioning as abrasion barriers, absorbing waste, and providing environmental tourism services (Dahuri *et al.*, 2004; Ratini, 2016) <sup>[12]</sup>. Coastal communities have long utilized mangrove plant parts for medicinal purposes, food, building materials, preservatives and coloring fishing nets commonly used by fishermen (Djamaludin, 2018).

Mangroves are almost exclusively distributed in the tropics (Hogarth, 2015; Ikhsanudin, 2018) <sup>[6]</sup>. This can be seen in the distribution of mangroves in the world, most of which are in tropical and warm climate countries. Indonesia with its geographical conditions, oceanographic characteristics and archipelago typology is very distinctive as a country that has high mangrove diversity and the largest area in the world (Djamaludin, 2018). Because of their high capacity for adaptation, mangrove communities show a high level ecological stability even though the physical and geological condition of the intertidal zone are prone to experiencing dramatic change (Alongi, 2015; Rahim, *et al.*, 2023) <sup>[11]</sup>.

Dulangeya Village is one of the villages in Botumoito District, Boalemo Regency, Gorontalo Province that has a lot of natural potentials that can be used for ecotourism, one of which is a quite extensive mangrove ecosystem. Mangrove ecosystems in Dulangeya village are often used by the surrounding community as building materials and also as ecotourism. Until now, there has been no study of the mangrove ecosystem in Dulangeya village.


This study aims to analyze the community structure of mangrove ecosystem constituents in Dulangeya Village which includes the number, species, density, frequency, and importance value index.

---

**Research Methods**

**Research Time and Site**

This research was conducted for 3 months from January to March 2023 in Dulangeya Village, Botumoito Sub-district, Gorontalo Province. Dulangeya Village is one of the coastal villages located in Boalemo Regency, Gorontalo Province with geographical boundaries of 0.48846°N, 122.24211°BT. The research location map is presented in Figure 1.



**Fig 1:** The research location map

**Tools and Materials**

The tools and materials and their uses used in this study are presented in Table 1.

**Table 1:** Tools and materials and their uses

Tools and materials	Uses
Mangrove identification book	For mangrove identification
Thermometer	For measuring temperature
Refraktometer	Salinity measurement
pH meter	pH measurement

**Sampling Technic Method**

The method used in this research is direct observation. Sampling for mangrove vegetation was done with a 100m long line transect or line transect as many as 4 stations. Determination of the station is based on the type of activity or activity that occurs around the station point. Stations 1 and 2 are adjacent to the boat mooring which is used as a tourist site, station 3 is adjacent to the house, and station 4 is adjacent to the pond. Each transect consists of plots with a size of 10x10m. Then identified each type of mangrove found in each plot based on mangrove identification book.

**Data Analysis**

To find out the community structure of mangrove vegetation, the parameters measured were mangrove density, relative density, frequency, relative frequency, species coverage, relative coverage, and important value index. They were using the following formula:

$$\text{Species density (ind/ m}^2\text{)} = \frac{\text{number of individuals}}{\text{total sampled areas}}$$

$$\text{Relative density (\%)} = \frac{\text{number of individuals in each species}}{\text{total individuals of all species}}$$

$$\text{Species frequency} = \frac{\text{number of plots in which a species occurs}}{\text{total sampled plots}}$$

$$\text{Relative frequency} = \frac{\text{frequency of a species}}{\text{several frequencies for all species}}$$

$$\text{Important value index} = \text{relative density} + \text{relative frequency} + \text{relative coverage}$$

**Result**

**Mangrove Environmental Conditions**

Based on the measurement results, the water conditions in Dulangeya Village are generally still favorable for mangrove growth. Water conditions in Dulangeya Village are presented in Table 2.

**Table 2:** Water condition of Dulangeya Village

No.	Parameters of oceanography	Value			
		St. 1	St. 2	St. 3	St. 4
1	pH	7.6	7.5	8.1	7.95
2	Temperature	28.8	27.4	30.6	30.55
3	Salinity	32	29.3	30	30

Based on the Decree of the Minister of Environment No. 51 of 2004 concerning seawater quality criteria for mangroves, a good pH ranges from 7-8.5, temperature ranges from 280C-320C, and salinity up to 34.

**pH**

The measurement results show that the pH value in the waters of Dulangeya Village ranges from 7.5 - 8.1. Siringoringo *et al*, (2018) [15] have measured pH levels in the Ciasem Protected Forest Area and obtained results ranging from 6.7 - 7.6. Each water area has a different pH value because it is influenced by the oceanographic and geomorphological characteristics of the area (Schaduw, 2018) [13].

**Temperature**

The measurement results showed that the temperature value ranged from 27.40C - 30.60C. The appropriate temperature for mangrove growth is more than 200C in the tropics (Aksornkoae, 1993; Schaduw, 2018) [13]. Water temperature measurements in the coastal mangrove forest of Amahai, Central Maluku Regency, ranged from 290C - 300C (Wailisa *et al*, 2022) [17]. Darwati *et al* (2022) [2] stated that differences in water temperature in an area can occur due to season, the circular position of the sun, and wind movement/circulation. In addition, the temperature affects physiological periods such as photosynthesis and evaporation, also affects salinity around mangroves (Maulidia, *et al*, 2022) [8].

**Salinity**

The salinity value in the waters of Dulangeya Village ranges from 29.3‰ - 32‰. This value is different from the results of salinity measurements in the waters of Kampung Sejahtera Bengkulu City which have values ranging from 16‰ - 20‰ (Putri *et al*, 2021) [10]. Pratiwi *et al* (2022) [9] stated that the lower the salinity value obtained, the closer to the river and the higher the salinity value obtained, the farther from the river area.

**Mangrove Vegetation**

The composition of mangrove species found in Dulangeya Village is 7 species, namely *Rhizophora stylosa*, *Rhizophora mucronata*, *Rhizophora apiculata*, *Brugueira gymnoriza*, *Brugueira cylindrica*, *Ceriops decandra*, *Ceriops tagal*,

*Scyphiphora hydrophyllacea*.**Station 1**

Mangrove species found at station 1 are 5 species with different values of relative density (RD), relative frequency (RF), and importance value index (IVI). (Table 3).

**Table 3:** Mangrove vegetation of station 1

Categories	Name of species	RD (%)	RF (%)	IVI (%)
Seedlings	<i>Rhizophora stylosa</i>	2.17	3.33	5.50
	<i>Rhizophora mucronata</i>	1.44	6.67	8.11
	<i>Rhizophora apiculata</i>	47.65	30.00	77.65
	<i>Brugueira gymmoriza</i>	19.86	30.00	49.86
	<i>Ceriops decandra</i>	28.88	30.00	58.88
Saplings	<i>Rhizophora stylosa</i>	5.21	11.76	16.97
	<i>Rhizophora mucronata</i>	1.67	2.94	4.61
	<i>Rhizophora apiculata</i>	11.46	11.76	23.22
	<i>Brugueira gymmoriza</i>	17.71	32.35	50.06
	<i>Ceriops decandra</i>	63.96	41.18	105.13
Tree	<i>Rhizophora stylosa</i>	11.21	15.09	30.89
	<i>Rhizophora mucronata</i>	13.55	16.98	66.11
	<i>Rhizophora apiculata</i>	50.93	30.19	127.16
	<i>Brugueira gymmoriza</i>	13.55	20.75	43.87
	<i>Ceriops decandra</i>	10.75	16.98	31.97

For the seedling category, the highest relative density value is owned by *Rhizophora apiculata*, the relative frequency value for *Rhizophora apiculata*, *Brugueira gymmoriza* and *Ceriops decandra* species has the same high value. The highest importance value index is owned by *Rhizophora apiculata*.

The sapling category for relative density, relative frequency and importance index with the highest value is owned by *Ceriops decandra*. Furthermore, for the tree category, *Rhizophora apiculata* has the highest value for relative density, relative frequency, and important value index. This shows that station 1 is dominated by *Rhizophora apiculata*.

**Station 2**

Mangrove species found at station 2 are 7 species with different values of relative density, relative frequency, and importance value index (Table 4).

**Table 4:** Mangrove vegetation of station 2

Categories	Name of species	RD (%)	RF (%)	IVI (%)
Seedlings	<i>Rhizophora stylosa</i>	37.80	25.00	62.80
	<i>Rhizophora mucronata</i>	13.72	20.83	34.55
	<i>Rhizophora apiculata</i>	0.61	4.17	4.78
	<i>Brugueira gymmoriza</i>	13.41	20.83	34.25
	<i>Ceriops decandra</i>	34.45	29.17	63.62
Saplings	<i>Rhizophora stylosa</i>	1.41	9.52	10.94
	<i>Rhizophora mucronata</i>	1.41	4.76	6.18
	<i>Brugueira gymmoriza</i>	31.80	28.57	60.37
	<i>Brugueira cilindrica</i>	1.77	4.76	6.53
	<i>Ceriops decandra</i>	57.24	38.10	95.34
Tree	<i>Ceriops tagal</i>	6.36	14.29	20.65
	<i>Rhizophora stylosa</i>	31.30	20.00	85.85
	<i>Rhizophora mucronata</i>	15.65	16.67	66.44
	<i>Rhizophora apiculata</i>	19.13	23.33	64.73
	<i>Brugueira gymmoriza</i>	10.43	13.33	25.77
Tree	<i>Ceriops decandra</i>	14.78	16.67	36.54
	<i>Ceriops tagal</i>	8.70	10.00	20.66

*Rhizophora stylosa* has the highest value for relative density in the seedling category, while *Ceriops decandra* dominates with the highest relative frequency and importance index values for the seedling category.

The saplings for the *Ceriops decandra* species dominate at station 2 with the highest relative density, relative frequency, and importance index values among other species. As for the tree category, the highest relative frequency value is owned by *Rhizophora apiculata*, and *Rhizophora stylosa* which has the highest relative density value and importance value index at station 2.

**Station 3**

Mangrove species found at station 3 are 7 species with different values of relative density, relative frequency, and importance value index (Table 5).

**Table 5:** Mangrove vegetation of stasiun 3

Categories	Name of species	RD (%)	RF (%)	IVI (%)
Seedlings	<i>Rhizophora stylosa</i>	18.60	7.41	26.00
	<i>Rhizophora mucronata</i>	9.09	18.52	27.61
	<i>Rhizophora apiculata</i>	7.85	11.11	18.96
	<i>Brugueira cilindrica</i>	14.46	18.52	32.98
	<i>Ceriops decandra</i>	6.20	11.11	17.31
	<i>Ceriops tagal</i>	34.30	18.52	52.82
Saplings	<i>Scyphiphora hydrophyllacea</i>	9.50	14.81	24.32
	<i>Rhizophora stylosa</i>	1.95	8.00	9.95
	<i>Rhizophora mucronata</i>	11.72	12.00	23.72
	<i>Rhizophora apiculata</i>	19.14	20.00	39.14
	<i>Brugueira cilindrica</i>	5.86	16.00	21.86
	<i>Ceriops decandra</i>	7.03	8.00	15.03
Tree	<i>Ceriops tagal</i>	50.39	28.00	78.39
	<i>Scyphiphora hydrophyllacea</i>	3.91	8.00	11.91
	<i>Rhizophora stylosa</i>	15.96	11.54	41.41
	<i>Rhizophora mucronata</i>	19.15	19.23	70.26
	<i>Rhizophora apiculata</i>	40.43	30.77	109.96
	<i>Brugueira cilindrica</i>	9.57	15.38	33.64
Tree	<i>Ceriops decandra</i>	1.06	3.85	5.25
	<i>Ceriops tagal</i>	13.83	19.23	39.49

The seedling category at station 3 for the same highest relative frequency value is owned by 3 species namely *Rhizophora mucronata*, *Brugueira cilindrica* and *Ceriops tagal*. In addition, *Ceriops tagal* also has the highest value for relative density and importance index.

*Ceriops tagal* dominates the sapling category by having the highest values for relative density, relative frequency, and important value index at station 3. Meanwhile, the tree category is dominated by *Rhizophora apiculata* which has the highest values for relative density, relative frequency, and important value index.

**Station 4**

Mangrove species found at station 4 are 5 species with different values of relative density, relative frequency, and importance value index (Table 6).



**Table 6:** Mangrove vegetation of station 4

Categories	Name of species	RD (%)	RF (%)	IVI (%)
Seedlings	<i>Rhizophora stylosa</i>	8.93	10.00	18.93
	<i>Rhizophora mucronata</i>	0.89	10.00	10.89
	<i>Rhizophora apiculata</i>	22.32	20.00	42.32
	<i>Brugueira gymnoriza</i>	6.25	20.00	26.25
Saplings	<i>Ceriops decandra</i>	61.61	40.00	101.61
	<i>Rhizophora stylosa</i>	16.67	26.09	42.75
	<i>Rhizophora apiculata</i>	16.67	17.39	34.06
	<i>Brugueira gymnoriza</i>	7.87	21.74	29.61
Tree	<i>Ceriops decandra</i>	58.80	34.78	93.58
	<i>Rhizophora stylosa</i>	14.50	17.14	39.18
	<i>Rhizophora mucronata</i>	7.63	17.14	41.76
	<i>Rhizophora apiculata</i>	43.51	25.71	112.37
	<i>Brugueira gymnoriza</i>	12.98	20.00	58.17
	<i>Ceriops decandra</i>	21.37	20.00	48.51

*Ceriops decandra* dominates for seedling and sapling categories at station 4, which has the highest relative density, relative frequency, and importance index values among other species. As for the tree category, it is dominated by *Rhizophora apiculata* with the highest relative density, relative frequency, and importance index among other species.

*Rhizophora* sp. is the most dominating species among other species, this type is most often found in all research stations ranging from seedling, sapling, to tree categories. This is in accordance with the statement of Usman *et al* (2013), that *Rhizophora* sp. is one of the mangrove species that is tolerant of environmental conditions (substrate, salinity, depth and nutrient sources) and can spread and grow in various places (Serosero *et al*, 2020)<sup>[14]</sup>. The type of substrate in Dulangeya Village is mud so it is very suitable for the habitat of *Rhizophora* sp. Muddy substrates in Indonesia are very suitable for the life of *Rhizophora* mangrove species (Rusila *et al*, 1999; Susanto *et al* 2018)<sup>[16]</sup>.

### Mangrove Damage

The standard criteria for mangrove damage determined through mangrove density values based on the Decree of the Minister of Environment No. 201 Year 2004 has been presented in Table 7.

**Table 7:** Result Mangrove damage standard criteria

Categories	Number of individuals (ind/ha)				Average	Criteria
	St. 1	St. 2	St. 3	St. 4		
Tree	1258.824	958.333	940	1091.667	1062.206	Moderate
Saplings	2823.529	2358.333	2560	1800	2385.466	Good
Seedlings	1629.412	2733.333	2420	933.333	1929.02	Good

The results showed that the mangrove ecosystem in Dulangeya Village was included in the moderate to good criteria. This is due to mangrove planting activities organized by the government and involving the village community so that the condition of the mangrove ecosystem in Dulangeya Village is still well maintained.

### Discussion

Based on the results of the study, 8 species of mangroves were found in four research locations in Dulangeya Village. They were *Rhizophora stylosa*, *Rhizophora mucronata*, *Rhizophora apiculata*, *Ceriops tagal*, *Ceriops decandra*, *Brugueira cylindrical*, *Brugueira gymnoriza*, and *Scyphiphora hydrophyllacea*. Species distribution of

mangrove has been influenced by many factors such as salinity, substrate type, and habitat geomorphology (Dharmawan and Pramudji, 2019).

The most common mangrove species with the highest importance index is *Rhizophora apiculata*. This shows that *Rhizophora apiculata* is a species that has the highest value of influence on the lives of coastal communities in Dulangeya Village. The large number of *Rhizophora apiculata* species found in a location is due to its high tolerance to extreme environmental conditions (Susanto *et al* 2018; Rahim *et al*, 2023)<sup>[11]</sup>. In addition, *Rhizophora apiculata* is also supported by a root system called aerial roots (pneumatophore), which can keep it stable (Hariyanto *et al*, 2019).

Based on the Decree of the Minister of Environment No. 201 Year 2004, mangrove ecosystem in Dulangeya Village is in good condition. This is due to the awareness of the surrounding community about the rules prohibiting mangrove logging.

### Conclusion

Based on the results of the study, there are 7 types of mangroves found in Dulangeya Village. Tree level density is classified as moderate and for sapling and seedling levels is classified as good. The highest species frequency for tree level is *Rhizophora apiculata*, while for sapling and seedling level is *Ceriops decandra*. The Index of Importance (INP) for the tree level obtained values ranging from 5.247 - 127.159. Mangrove ecosystems in Dulangeya Village are included in moderate to good criteria based on the standard criteria for mangrove damage determined by the Minister of Environment Decree No. 201 of 2004.

### Acknowledgement

The authors would like to thank the Dulangeya Village government for allowing the author to conduct research at that location. The authors also thank to the youth organization for helping the author in collecting data in the field.

### References

- Bengen DG. Synopsis of Coastal Ecosystems and Natural Resources. Center for Coastal and Marine Resources Studies. Bogor Agricultural Institute. 2000; 62 pages.
- Darwati H, E Poedjirahajoe, R Sadono, HB Soewarno. Water Characteristics and Dominant Mangrove Species in Panjang Island, Padang Tikar II Village, Kubu Raya Regency, West Kalimantan. Jurnal Hutan Lestari. 2022; 10(4):1002-1009.
- Dharmawan AWE, Pramudji. Mangrove Community Structure in Papua Small Islands, Case Study in Biak Regency. IOP: International Conference of Mangrove and Its Related Ecosystem. 2019, 1-8 IOP Conf. Ser.: Earth Environ. Sci. 550 012002.
- Djamaluddin R. Mangrove (Biology, Ecology, Rehabilitation, and Conservation). Unsrat Press. Manado, 2018.
- Hariyanto S, AK Fahmi, T Soedarti, EE Suwarni. Vegetation and Community Structure of Mangrove in Bama Resort Baluran National Park Situbondo East Java. Biosaintifika: Journal of Biology & Biology Education. 2019; 11(1):132-138.
- Ikhsanudin N. Utilization and Stakeholders of Mangrove Forest in Indramayu Regency, West Java Province. Graduate School, Bogor Agricultural University. 2018;

- 67 pages.
7. Karimah. The Role of Mangrove Ecosystems as Habitat for Marine Organisms. *Jurnal Biologi Tropis*, 2017; 17(2):51-58.
  8. Maulidia V, AA Akbar, Jumiati, Arifin, A Sulastri. The Value of Mangrove Ecosystem Based on Mangrove Carbon Sequestration in West Kalimantan. *Journal of Wetlands Environmental Management*. 2022; 10(1):12-26.
  9. Pratiwi FKWN, L Masluka, DN Sugianto. Water and Sediment Quality at the Mangrove Information Center (MIC), Pekalongan. *Indonesia Journal of Oceanography*. 2022; 4(3):33-43.
  10. Putri YD, E Yuliza, Lizalidiawati. Assessment of Water Quality and Pollution Index in the Waters of Prosperous Village of Baai Island, Bengkulu City. *Newton-Maxwell Journal of Physics*. 2021; 2(2):45-53.
  11. Rahim A, TR Soeprbowati, TT Putranto. Assessment of Mangrove Biodiversity and Community Structure as a Basis for Sustainable Conservation and Management Plan in Tambakbulusan, Demak, Central Java. *AACL Bioflux*. 2023; 16(2):753-767.
  12. Ratini. Mangrove Ecosystem Conservation Planning in Ujung Alang Village, Kampung Laut District, Cilacap Regency. Graduate School, Bogor Agricultural University. 2016; 87.
  13. Schaduw JNW. Community structure and sustainability of mangrove ecosystem management on small islands (case of Nain Island, North Minahasa Regency, North Sulawesi Province). *Jurnal Ilmu Lingkungan*. 2018; 16(2):120-129.
  14. Serosero RH, S Abubakar, S Hasan. Distribution and Community Structure of Mangrove in Donrotu, Guratu and Manomadehe Islands, West Halmahera District, North Maluku. *Jurnal Ilmu dan Teknologi Kelautan Tropis*. 2020; 12(1):151-166.
  15. Siringoringo HH, BH Narendra, AG Salim. Water Quality of Mangrove Ciasem Pamanukan Subang Regency, West Java. *Jurnal Pengelolaan Sumberdaya Alam dan Lingkungan*. 2018; 18(3):301-307.
  16. Susanto AH, T Soedarti, H Purnobasuki. Mangrove Community Structure of Surabaya East Coast. *An International Journal of Plant Research & Biotechnology*. 2018; 31(3):09-12.
  17. Wailisa R JD Putuhena, F Soselisa. Analysis of Water Quality in the Coastal Mangrove Forest of Negeri Amahai, Central Maluku Regency. *Jurnal Hutan Pulau-Pulau Kecil*. 2022; 6(1):57-71.