



Potential of *Tephrosia candida* for Restoration of Degraded Lands and Fuel Wood Production in Western Ethiopia

Mezgebu Senbeto Duguma ^{1*}, Regassa Terefe Urga ²

¹ Researcher at Bako Agricultural Research Center, Oromia Agricultural Research Institute (IQQO), Ethiopia

² Project Manager at the Hunger Project, Ethiopia

* Corresponding Author: **Mezgebu Senbeto Duguma**

Article Info

ISSN (online): 2582-7138

Volume: 04

Issue: 04

July-August 2023

Received: 24-05-2023

Accepted: 19-06-2023

Page No: 318-323

Abstract

The challenges in restoration projects are to find potential and compatible tree and shrub species to start and scale up them in a way that will have a clear effect at the landscape level while these species can help to achieve climate-smart agriculture's "triple wins" of increasing rural incomes and biomass energy supply. This study aimed to examine the potential of *Tephrosia candida* as a species for degraded farmland restoration thereby to evaluate its biomass for fuel wood and local energy source. There are 3 locations where each location has 6 plots. Each plot contains 25 plants with 2m*2m spacing. Composite soil sample, plant height, DBH, survival rate, dry weight, crown cover, number of branches and time of seed collection were collected. Growth performance of *Tephrosia candida* was very fast which showed about 15cm height and 0.26cm diameter increment monthly, and its survival was 100% at all sites. Averagely 1.5 branches/Month recorded while crown cover increased accordingly, and on average 25 main branches and 38 secondary branches, and each shrub weigh 22.22kg on average when dried. Each *Tephrosia*'s crown cover was 16.25m² on average. Analysis of composite soil sample and after 3 years of implementation soil pH increment observed in all locations, accordingly, soil pH at Diga improved from 4.89 (before) to 5.22 (3 years after implementation). We recommend *Tephrosia candida* for fast restoration purposes and farmland fuel wood producing species to cover the increasing fuel wood demand due to its high potential of both restoration and high biomass production.

Keywords: Biomass energy, fuel wood, restoration, Shrub species *Tephrosia candida*

1. Introduction

1.1 Restoration and its limitations

In most developing and developed countries, the issues of energy, food and shelter demands and supplies are mainly depended directly or indirectly on natural resources, specially, on forest and land resources which is especially true for poor and vulnerable rural communities where land is often the most important asset, and land itself and other natural resources are under risk of degradations (IPBES, 2018; FAO and UNCCD 2022) ^[25, 12]. On the other hand, rapidly increasing population, deforestation, growing food waste and declining soil fertility are posing serious challenges to humanity for future food security where the land degradation must be solved (Gupta, 2019) ^[18]. As a result, currently, degradation of the Earth's land surface through human activities is negatively impacting the well-being of at least 3.2 billion people, loss of biodiversity and ecosystem services (FAO 2021, IPBES 2018) ^[25]. Achieving restoration of degraded lands can provide multiple benefits in terms of improving livelihoods, increasing bioenergy sources, empowering women and youth, and enhancing both biodiversity conservation and climate change mitigation and adaptation (FAO and UNCCD 2022, FAO 2018) ^[12].

The challenges in restoration projects are to find potential and compatible tree and shrub species to start and scale up them in a way that will have a clear effect at the landscape level while these species can help to achieve climate-smart agriculture's "triple wins" of increasing rural incomes, biomass energy supply, making agricultural products more resilient in the face of climate extremes, and making agriculture a solution to the climate change problem rather than part of the problem (Deweese *et al.*, 2011) [8].

Africa is one of the most vulnerable continents in the world. It has over 700 million ha of its land already degraded, and degradation still occurs at an alarming rate of 3% annually. This offers tremendous opportunity for restoration. In the last two centuries, humans have converted or modified 70% of the world's grasslands, 50% of the savannah, 45% of the temperate deciduous forest, and 27% of the tropical forest biome primarily for farming and grazing activities (FAO 2021). In twenty years (1985-2005), the world's croplands and pastures expanded by 154 million hectares (Ray *et al.*, 2012) [39]. This expansion has dramatically increased food production (Mansourian *et al.* 2015) [43, 34], but at the expense of many life-supporting ecosystem services on which our wellbeing, and that of future generations, depend (Duguma *et al.* 2019) [35]. Agriculture and fuel wood are estimated to be the proximate driver for more than 80% of deforestation and land degradation worldwide (Kissinger and Desy, 2012) [30], which are likely to continue in order to meet the projected increases in demand for food, energy and water in the next decades (FAO 2021, FAO 2018). In 2016, Ethiopia faced the worst drought in fifty years that affected more than 10 million people (FAO, 2018). Although some articles discuss that over the last 20 years, the scale of restoration of degraded land through soil and land conservation and tree planting in Ethiopia is noteworthy, there are still tremendous land degradation continues in most parts of the country in general, and specifically in western parts of Ethiopia.

Tephrosia is a genus of leguminous shrubby plants and herbs mostly found in tropical countries (Blommaert, 1950). The plant is established easily from seeds (Barnes and Freyre, 1969) [1]. The foliage is often ashy-pubescent, hence the name *Tephrosia* (Gadzirayi *et al.* 2009) [16]. *Tephrosia* can be grown to improve soil fertility, for firewood, as an insecticide against storage pests and mites on plants. It is best species in open grazing areas, because goats and other animals usually do not like to browse the trees, and it is also quite resistant to termites (Garrity *et al.* 2010; Munthali *et al.* 2014) [17, 36].

1.2. Biomass energy demands and natural resources management

In sub-Saharan Africa countries, over two-thirds of the households use wood as their main fuel for cooking, heating and water boiling. In Africa, demand for wood is likely to continue where in Sub Saharan Africa, fuel wood dominates the use of wood, as it is the most important energy source (FAO 2017, De Miranda *et al.* 2013) [6], and the fuel wood consumption in Africa is predicted to increase in the consecutive decades from 2000 (FAO 2017; Sepp *et al.* 2014; Hoffmann *et al.* 2016; Sander *et al.* 2011; Dewees *et al.*, 2011) [8, 42, 22, 41].

Whatever lands are chosen, bioenergy can strengthen the economic case for restoration (IRENA 2017) [27] while the worldwide use of bioenergy has been increasing greatly in recent years (IEA Bioenergy, 2016; IRENA, 2016) [24, 26], which was mainly driven by an increase in demand for low-

carbon energy. For this purpose, there are different potential fuel wood plants, native and nonnative, which are acquired from the forest reserve or from people's homes and farmlands (Kaburi and Medley 2011) [29], while these plants differ by their material uses, attributes as a fuelwood, and ease of propagation (Fischer *et al.*, 2012) [14]. Therefore, in our study, an on-farm experiment was established in 2020 in western Oromia, Ethiopia to examine the potential of *Tephrosia candida* as a species for degraded farmland restoration thereby to evaluate its biomass for fuel wood and local energy source as it can minimize burdens on natural forest resources in the 3 locations of the study area.

2. Materials and Methods

An on-farm experiment was established in 2020 in western Oromia, Ethiopia, and the three locations were Bako, Diga and Wayu Tuqa districts of western Oromia. The selected locations are where soil acidity is very high and land degradation is very severe. From different species of *Tephrosia*, *Tephrosia candida* was used here. We planted 150 *Tephrosia* plants in each location where there were 6 plots on each. Each plot contains 25 plants with 2m * 2m spacing.

All data including initial composite soil sample, growth parameters (height, DBH, survival rate), dry weight, morphological expressions (crown cover, number of branches) were collected, and an average data from each location were taken for further analysis.

3. Results and Discussion

3.1. Growth performance, dry weight and crown cover of *Tephrosia candida*

Data collection and measurement of its growth performances starting from germination were carefully conducted. The growth performance of *Tephrosia candida* was very fast which showed about 15cm height and 0.26cm diameter increment monthly, and its survival was 100% at all sites. Emergences of branches were averagely 1.5 branches per Month while crown cover increased accordingly. This performance of the tephrosia shrub will make it very potential for fuel wood and fast facilitation of restoration which also agree with Mafongoya and his colleagues finding, which says, at three months after planting, tephrosia exhibited good initial growth with the fastest growth with 138 cm, and wood biomass yield produced after nine months of growth was highest among other shrub species for *Tephrosia candida* (Mafongoya *et al.*, 2003) [31]. Due to its fast growing and many branches bearing in short duration of time, tephrosia is preferably recommended for the purposes of restoration and bioenergy production at farmers level in which it can be used as an improved fallow agroforestry practices. This issue was briefly discussed in Christersson *et al.*, (2006) [5], and in their finding they pointed that high biomass production, which could be influenced by man, include the choice of tree species, provenances and clones, together with good resistance to insects, fungus, virus and browsing animals and hardness to frost and drought, where it is important to use trees species, proveniences and clones that have the potential for a rapid and dense canopy development where a design of the plantation is of great importance, and we recommend for such demands to use the *Tephrosia candida* shrub. Additionally, its fast growing habit is better than other leguminous and nitrogen fixing shrubs that it is also concluded better than *Leucaena leucocephala* and *Gliricidia sepium* species in Odedire and Babayemi (2007) [38] and

Makumba and Phiri (2008) ^[32] studies respectively.

A data collected after 3 years of planting showed that *Tephrosia* could bear on average 25 main branches and 38 secondary branches, and each *Tephrosia* shrub can weigh 22.22kg on average when dried. And also, each *Tephrosia*'s crown biomass can cover the area of 16.25m² on average (Tab. 1). In line with different studies and findings of many researchers, because of high demand and high consumption degrees of fuel wood in Africa, especially, in the sub-Saharan Africa countries, and also to fulfill the need of fast and high biomass bearing tree and/or shrub species, using *tephrosia* species is highly recommended by the current finding. This result can also contribute to Van Loon 2017 ^[46], IUCN 2016 and the Bonn Challenge findings and worries, which were highly described that the degraded lands are challenging and economically unattractive for food crop cultivation, biomass plantations with high-yielding wood or grass species can

allow bioenergy and also can help to restore and reclaim such lands to be extracted without conflicting with food production thereby improve carbon storage. Additionally, growing wood on degraded land can further serve to curb unsustainable wood extraction from local forest resources (Van Loon, 2017) ^[46]. The best initiative of the Bonn Challenge, for example, provides a great opportunity to unlock the potential of degraded land which put its goal to a global endeavor to restore a total of 350 million hectares (Mha) by 2030, where 150 Mha projected by 2020, and another 200 Mha by 2030. Accordingly, 18 sub-Sahara African countries have pledged to restore some 75 Mha through AFR100, where the African Forest Restoration initiative aiming to regenerate 100 Mha (<http://www.bonnchallenge.org/>, Van Loon, 2017, IUCN 2016) ^[46].

Table 1: Morphological Expression of *Tephrosia candida*

Location	Plot No	Dry weight (kg)	Crown cover			Branch No	
			Radius(m)	R2(r*r)	Area(m2)=3.14* R2	Main	2ndary
Bako	1	22	2.3	5.29	16.61	26	37
Bako	2	21	2.2	4.84	15.20	25	38
Bako	3	24	2.5	6.25	19.63	27	38
Bako	4	23	2.3	5.29	16.61	24	36
Bako	5	22	2.3	5.29	16.61	26	39
Bako	6	21.5	2.1	4.41	13.85	25	37
Average		22.3	2.3	5.2	16.4	25.5	37.5
Diga	1	21	2.1	4.41	13.85	23	35
Diga	2	22.5	2.3	5.29	16.61	22	36
Diga	3	21	2.1	4.41	13.85	25	38
Diga	4	20	2.1	4.41	13.85	24	35
Diga	5	21	2	4	12.56	22	36
Diga	6	19	2.2	4.84	15.20	23	34
Average		20.75	2.13	4.56	14.32	23.17	35.67
Wayu Tuqa	1	19.5	2.5	6.25	19.63	24	37
Wayu Tuqa	2	21	2.35	5.5225	17.34	26	36
Wayu Tuqa	3	20	2.3	5.29	16.61	26	37
Wayu Tuqa	4	23	2.4	5.76	18.09	24	35
Wayu Tuqa	5	21	2.3	5.29	16.61	25	37
Wayu Tuqa	6	20.5	2.1	4.41	13.85	24	38
Average		20.83	2.33	5.42	17.02	24.83	36.67

3.2. Comparison of Growth Performance at the 3 Locations

Tephrosia candida shows well performance and 100%

survival at all locations. There were no significant difference among dry weight biomass, crown cover, and sprouting of main and secondary branches (Fig. 1).




Fig 1: Graph of growth performances in all the 3 locations

From this graph, we can conclude that tephrosia can be well grown uniformly in different acidic and structurally different soil type areas, because these locations have different soil structures, fertility status and soil acidity levels.

3.3. Soil acidity improvement

Tephrosia candida is a shrub species with crowded crown and a shallow root which has very fast potential or ability to soften a compacted soil. Soil features under old tephrosia shrub is very soft, composted and looks better for production (Fig. 2). Composite soil sample (before tephrosia planting) was taken and pH values were analyzed. We also collected and analyze soil sample after 3 years of implementation, and its result shows that there is soil pH increment in all locations, for example, soil pH at Diga improved from 4.89 (before) to 5.22 (3 years after implementation) (Tab. 2). This result agrees with the findings of many studies who were explaining that tephrosia can be grown to improve soil fertility, for firewood, as an insecticide against storage pests and mites on plants, and it is also quite resistant to termites (Garrity *et al.* 2010; Munthali *et al.* 2014) ^[17, 36].

Table 2: Soil pH values before and after tephrosia plantation

Location	Soil pH before implementation	Soil pH after 3 years
Bako	5.11	5.47
Diga	4.89	5.22
Wayu Tuqa	5.16	5.51

From the table, we can understand that restoring degraded and acidic soil with tephrosia shrubs, and using them as an improved fallow can improve soil acidity problems thereby can improve vegetation survival even on the highly acidic soils within short period of improved fallow times. This was also discussed in the findings of Garrity *et al.* (2010) ^[17] and Munthali *et al.* (2014) ^[36] that tephrosia roots, leaves and seeds contain high amounts of nutrients, especially nitrogen, which is necessary for good plant development, and when the trees are cut and the leaves worked into the soil, these nutrients can be used by the plants that are grown after Tephrosia in the field.



Fig 2: Photo of old and failed *Tephrosia candida*, taken by the first author

From the picture, we can observe that whatever the soil from the beginning, tephrosia will soften the soil after few years and might be left the land for other agricultural activities so that it will be failed and used as a fuel wood.

3.4. Seed Collection and Handling

Tephrosia can in many ways be used just like pigeon pea, and seed harvest, storage and sowing are done in the same way. After collection of brown pods and threshing it, it is better to dry the seeds for at least 3 to 4 days under shade. In western Oromia, peak seed production is between November and February. However, it can bear to produce smaller quantities of seed throughout the rest of the year.

4. Conclusion and Recommendations

Over the last 20 years, the scale of restoration of degraded land through soil and land conservation and tree planting in Ethiopia is remarkable, however, there are still tremendous land degradation continues in most parts of the country in general, and specifically in western parts of Ethiopia. Potential species selection for restoration and multifunctional purposes is still challenging in most cases and many restoration projects are facing low survival rate of seedlings after planting. *Tephrosia candida* is a shrub species with crowded crown and a shallow root which has very fast potential or ability to soften a compacted soil even in acidity soils and high above ground biomass production. Restoring degraded and acidic soil with tephrosia shrubs, and using the shrub species as an improved fallow could also improve soil acidity problems and also can improve vegetation survival even on highly acidic soils within short period of times. Tephrosia is seen quite better in growth performance, survival rate, reclaiming soil acidity, dry biomass provision and softening compacted soils than other shrub species. Additionally, since it is not as palatable as other forage shrubs which attract goat, sheep, cattle and other browsing wild animals it is very potential for restoration activities, especially in open grazing countries like Ethiopia. Its seed can be collected and handled simply and it has seen that there is no germination problem for further production. Its fast growing performance and its short life cycle which support the shrub to rotate a farmland where it had been grown to any other crop production magnifies its demands. Therefore, this finding recommends *Tephrosia candida* shrub species for restoration of farmlands, and fuel wood production purposes within short period of time.

5. Funding: This research was funded by Oromia Agricultural Research Institute (IQQO).

6. Acknowledgments: My acknowledgement also goes for all Bako Agricultural Research Center (BARC) staffs, and also to all DAs and experts of Wayu Tuqa and Diga districts.

7. Conflicts of Interest: The author declares no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

8. References

1. Barnes DK, Freyre RH. Seed-production potential of *Tephrosia vogelii* in Puerto Rico. The Journal of Agriculture of the University of Puerto Rico. 1969.
2. Beisner BE, Haydon DT, Cuddington K. Alternative stable states in ecology. Frontiers in Ecology and the Environment. 2003;1(7):376-82.
3. Berhane G, Hoddinott JF, Kumar N, Taffesse AS, Diressie MT, Yohannes Y, *et al.* Evaluation of

- Ethiopia's Food Security Program; c2012.
4. Blommaert KLJ. The plant *Tephrosia vogelii* Hooker as a freshwater fish poison. Transactions of the Royal Society of South Africa. 1949;32(3):247-264.
 5. Christersson L, Verwijst T, Amatya SM. Wood production in agroforestry and in short-rotation forestry systems: Synergies for rural development; c2006.
 6. De Miranda RC, Sepp S, Ceccon E, Owen M. Commercial woodfuel production: Experience from three locally controlled wood production models. The World Bank. 2013;75195:1-52.
 7. Desta L, Carucci V, Wendem-Agenehu A, Abebe Y. Community-based participatory watershed development: A guideline. Annex; c2005.
 8. Dewees PF, Place SJ, Scherr C, Buss. Investing in trees and landscape restoration in Africa: What, where, and how. Washington, DC: Program on Forests (PROFOR); c2011.
 9. Woolf D, Solomon D, Lehmann J. Land restoration in food security programmes: Synergies with climate change mitigation. Climate Policy. 2018;18(10):1260-1270. DOI: 10.1080/14693062.2018.1427537.
 10. Fagerström MH, Nilsson SI, Van Noordwijk M, Phien T, Olsson M, Hansson A, *et al.* Does *Tephrosia candida* as fallow species, hedgerow, or mulch improve nutrient cycling and prevent nutrient losses by erosion on slopes in northern Vietnam? Agriculture, Ecosystems & Environment. 2002;90(3):291-304.
 11. FAO. The state of the world's land and water resources for food and agriculture (SoLaW): Managing systems at risk. Food and Agriculture Organization of the United Nations, Rome and Earthscan, London; c2011.
 12. FAO and UNCCD. Technical guide on the integration of the voluntary guidelines on the responsible governance of tenure of land, fisheries and forests in the context of national food security into the implementation of the United Nations Convention to combat desertification and land degradation neutrality. FAO, Rome and UNCCD, Bonn; c2022. <https://doi.org/10.4060/cb9656en>.
 13. FAO. The future of food and agriculture: Alternative pathways to 2050. Food and Agriculture Organization of the United Nations, Rome; c2018.
 14. Fischer J, Hartel T, Kuemmerle T. Conservation policy in traditional farming landscapes. Conservation Letters. 2012;5(3):167-175.
 15. Food and Agriculture Organisation. Incentivizing sustainable wood energy in sub-Saharan Africa: A way forward for policy-makers; c2017.
 16. Gadzirayi C, Mutandwa E, Mwale M, Chindundu T. Utilization of *Tephrosia vogelii* in controlling ticks in dairy cows by small-scale commercial farmers in Zimbabwe. African Journal of Biotechnology. 2009;8(17):4134-4136.
 17. Garrity DP, Akinnifesi FK, Ajayi OC, Weldesemayat SG, Mowo JG, Kalinganire A, *et al.* Evergreen agriculture: A robust approach to sustainable food security in Africa. Food Security. 2010;2:197-214.
 18. Gupta GS. Land degradation and challenges of food security. Review of European Studies. 2019;11:63.
 19. Hobbs RJ. The ecological context: A landscape perspective. In: Perrow MR, Davy AJ, editors. Handbook of Ecological Restoration, Vol. 1: Principles of Restoration. Cambridge: Cambridge University Press; c2002. p. 22-45.
 20. Hobbs RJ. Ecological filters, thresholds, and gradients in resistance to ecosystem reassembly. In: Temperton VM, Hobbs RJ, Nutton T, Halle S, editors. Assembly Rules and Restoration Ecology: Bridging the Gap Between Theory and Practice. Washington, D.C.: Island Press; c2004. p. 72-95.
 21. Hobbs RJ. Setting effective and realistic restoration goals: key directions for research. Restoration Ecology. 2007;15(2):354-357.
 22. Hoffmann H, Brüntrup M, Dewes C. Wood energy in sub-Saharan Africa: how to make a shadow business sustainable (No. 14/2016). Briefing Paper. c2016.
 23. Hooke RL, Martín-Duque JF, Pedraza J. Land transformation by humans: A review. Satoday. 2012;22(12):4-10.
 24. IEA Bioenergy. IEA Bioenergy Countries' Report: Bioenergy Policies and Status of Implementation. International Energy Agency, Paris; c2016.
 25. IPBES. Report of the Plenary of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services on the work of its sixth session. Medellín, Colombia, c2018, 18-24.
 26. IRENA. Renewable Energy Statistics. International Renewable Energy Agency, Abu Dhabi; c2016.
 27. IRENA. Bioenergy from degraded land in Africa: Sustainable and technical potential under Bonn Challenge pledges. International Renewable Energy Agency, Abu Dhabi; c2017.
 28. IUCN. Africa & the Bonn Challenge: A demonstration of leadership. IUCN Forest Brief; c2016.
 29. Kaburi SM, Medley KE. Community perspectives on fuelwood resources in East Africa. Mountain Research and Development. 2011;31(4):315-324.
 30. Kissinger G, Herold M, Desy V. Drivers of deforestation and forest degradation: A Synthesis Report for REDD+ policymakers. Lexeme Consulting, Vancouver, Canada; c2012.
 31. Mafongoya PL, Chintu R, Chirwa TS, Matibini J, Chikale S. *Tephrosia* species and provenances for improved fallows in southern Africa. Agroforestry Systems. 2003;59(3):279-288.
 32. Makumba W, Phiri A. Evaluation of *Gliricidia sepium* and *Tephrosia candida* leaves as a source of nitrogen for vegetable production in a biomass transfer. In: Annual Report-Soil and Agricultural Engineering Commodity Group, 2007/08. Presented during the DARS Annual Planning meeting held at Andrews Holiday Resort, Mangochi, 14-19th September. Department of Agricultural Research Services Headquarters. c2008;2-8.
 33. Mansourian S, Berrahmouni N. Review of forest and landscape restoration in Africa 2021. Food & Agriculture Organization; c2021.
 34. Mansourian S, Vira B, Wildburger C. Forests and food: addressing hunger and nutrition across sustainable landscapes. Open Book Publishers. c2015; p. 288.
 35. Senbeto DM, Feyssa DH, Biber-Freudenberger L. Agricultural biodiversity and ecosystem services of major farming systems: A case study in Yayo Coffee Forest Biosphere Reserve, Southwestern Ethiopia. MDPI Agriculture. Germany; c2019;1-26.
 36. Munthali MG, Gachene CK, Sileshi GW, Karanja NK. Amendment of *Tephrosia* improved fallows with inorganic fertilizers improves soil chemical properties,

- nitrogen uptake, and maize yield in Malawi. International Journal of Agronomy; c2014.
37. Nkonya E, Anderson W, Kato E, Koo J, Mirzabaev A, von Braun J, *et al.* Global cost of land degradation. In: Economics of Land Degradation and Improvement – A Global Assessment for Sustainable Development. Cham: Springer; c2016. p. 117-165.
 38. Odedire JA, Babayemi OJ. Preliminary study on *Tephrosia candida* as a forage alternative to *Leucaena leucocephala* for ruminant nutrition in Southwest Nigeria. Livestock Research for Rural Development. 2007;19(9):1-11.
 39. Ray DK, Ramankutty N, Mueller ND, West PC, Foley JA. Recent patterns of crop yield growth and stagnation. Nature Communications. 2012;3:1293.
 40. Safriel U. The assessment of global trends in land degradation. In: Mannava VK, Ndegwa N, editors. Climate and Land Degradation. Berlin: Springer; c2007. p. 1-38.
 41. Sander K, Haider SW, Hyseni B. Wood-based biomass energy development for sub-Saharan Africa: Issues and approaches. The World Bank. 2011;74545:1-64.
 42. Sepp S, Sepp C, Mundhenk M. Towards sustainable modern wood energy development: Stocktaking paper on successful initiatives in developing countries in the field of wood energy development. Deutsche Gesellschaft für Internationale Zusammenarbeit (GIZ) GmbH, Bonn. Society for Ecological Restoration International Science and Policy Working Group; c2004-2014.
 43. Stanturf JA, Kant P, Lillesø JPB, Mansourian S, Kleine M, Graudal L, Madsen P. Forest landscape restoration as a key component of climate change mitigation and adaptation. International Union of Forest Research Organizations (IUFRO), Vienna, Austria. c2015;34.
 44. Suding KN, Gross KL, Houseman GR. Alternative states and positive feedbacks in restoration ecology. Trends in Ecology and Evolution. 2004;19(1):46-53.
 45. Temperton VM, Hobbs RJ, Nuttle T, Halle S, editors. Assembly rules and restoration ecology – Bridging the gap between theory and practice. Washington, D.C.: Island Press; c2004.
 46. Van Loon TJ. Restoring degraded land: On the bioenergy potential from the Bonn Challenge pledges in sub-Saharan Africa (Master's thesis); c2017.
 47. Dr. Priya Mehra. Role of Soil Microbiota in Ecosystem Restoration and Functionality. Journal of Soil Future Research. 2022;3(2):26-31.