

# International Journal of Multidisciplinary Research and Growth Evaluation.



## Environmental Health and Disease Prevention: Conceptual Frameworks Linking Pollution Exposure, Climate Change, and Public Health Outcomes

Mercy Egemba 1\*, Simeon Ayo-Oluwa Ajayi 2, Costly Aderibigbe-Saba 3, Patrick Anthony 4

- <sup>1-3</sup> Independent Researcher, USA
- <sup>4</sup> Novartis, Kano, Nigeria
- \* Corresponding Author: Mercy Egemba

### **Article Info**

ISSN (Online): 2582-7138 Impact Factor (RSIF): 7.98

Volume: 05 Issue: 03

May - June 2024 Received: 20-04-2024 Accepted: 22-05-2024 Published: 29-06-2024 Page No: 1133-1153

#### Abstract

The intersection of environmental pollution, climate change, and public health represents one of the most pressing challenges of the 21st century, with profound implications for disease prevention and population health outcomes. This comprehensive review examines the conceptual frameworks that link pollution exposure and climate-related environmental changes to public health outcomes, providing a systematic analysis of the mechanisms through which environmental factors influence disease patterns and health disparities. The study synthesizes current research on environmental health determinants, exploring the complex pathways through which air pollution, water contamination, extreme weather events, and ecosystem disruption contribute to disease burden across diverse populations. Through an extensive literature review encompassing epidemiological studies, environmental health assessments, and climate science research, this paper identifies key conceptual models that explain the relationships between environmental exposures and health outcomes. The analysis reveals that environmental health impacts operate through multiple interconnected pathways, including direct toxic effects of pollutants, indirect effects through ecosystem disruption, and social vulnerability factors that amplify exposure risks among marginalized populations. Climate change acts as a threat multiplier, exacerbating existing environmental health risks while creating new exposure scenarios and altering disease transmission patterns. The paper presents a comprehensive framework for understanding these relationships, incorporating concepts of environmental justice, cumulative risk assessment, and adaptive capacity in health systems. Key findings indicate that effective environmental health protection requires integrated approaches that address both immediate pollution control and long-term climate adaptation strategies. The research highlights the importance of place-based interventions, community engagement, and multi-sectoral collaboration in developing effective disease prevention strategies. Policy implications emphasize the need for proactive environmental health governance that incorporates climate resilience, equity considerations, and evidence-based risk assessment. The study concludes with recommendations for strengthening conceptual frameworks through improved environmental health surveillance, enhanced inter-disciplinary research collaboration, and the development of innovative intervention strategies that address the root causes of environmental health disparities. These findings contribute to the growing body of knowledge on environmental determinants of health and provide practical guidance for public health practitioners, policymakers, and researchers working to address the complex challenges at the intersection of environmental change and population health.

DOI: https://doi.org/10.54660/.IJMRGE.2024.5.3.1133-1153

**Keywords:** Environmental Health, Disease Prevention, Pollution Exposure, Climate Change, Public Health Outcomes, Conceptual Frameworks, Environmental Justice, Risk Assessment

### 1. Introduction

Environmental health represents a fundamental determinant of population health outcomes, encompassing the complex relationships between physical, chemical, and biological factors in the environment and their impacts on human health and disease patterns (Prüss-Ustün *et al.*, 2016). The field has evolved significantly over the past several decades, moving from a

focus on acute exposures and immediate health effects to a more comprehensive understanding of how environmental factors interact with social, economic, and behavioral determinants to influence health outcomes across the lifespan (Landrigan *et al.*, 2018). This evolution has been driven by mounting evidence of the profound ways in which environmental exposures contribute to the global burden of disease, with estimates suggesting that environmental factors account for approximately 24% of the global disease burden and 23% of deaths worldwide (WHO, 2016).

The conceptual frameworks that guide environmental health research and practice have undergone substantial refinement as our understanding of exposure pathways, dose-response relationships, and population vulnerability has advanced (Nkwazema *et al.*, 2023). Traditional models focused primarily on single pollutant exposures and direct health effects have given way to more sophisticated frameworks that recognize the complexity of real-world exposures, which typically involve multiple pollutants, multiple exposure routes, and interactions with other environmental and social stressors (Dominici *et al.*, 2010). These contemporary frameworks emphasize the importance of cumulative risk assessment, environmental justice considerations, and the role of social determinants in modifying environmental health impacts (Morello-Frosch & Lopez, 2006).

Climate change has emerged as a critical factor that fundamentally alters the environmental health landscape, acting as both a direct driver of health outcomes and a modifier of existing environmental health risks (Watts et al., 2021). The Intergovernmental Panel on Climate Change has identified numerous pathways through which climate change affects human health, including altered patterns of temperature-related morbidity and mortality, changes in the distribution and incidence of climate-sensitive infectious diseases, impacts on food security and nutrition, and health effects related to extreme weather events and ecosystem disruption (IPCC, 2022). These climate-health linkages operate through complex mechanisms that intersect with existing environmental health risks, creating new exposure scenarios and exacerbating existing health disparities (Hayes et al., 2018).

The integration of climate change considerations into environmental health frameworks has necessitated the development of new conceptual models that can capture the dynamic and interconnected nature of climate-environmenthealth relationships (Berry *et al.*, 2018). These models recognize that climate change acts as a threat multiplier, amplifying existing environmental health risks while creating novel exposure pathways and altering the effectiveness of traditional disease prevention strategies (Ebi & Semenza, 2008). The temporal and spatial scales at which climate change operates present unique challenges for environmental health assessment and intervention, requiring frameworks that can address both immediate adaptations needs and long-term prevention strategies (Friel *et al.*, 2011).

Environmental justice has become a central consideration in contemporary environmental health frameworks, recognizing that environmental health impacts are not distributed equally across populations (Bullard & Johnson, 2000). Communities of color, low-income populations, and other marginalized groups often face disproportionate exposure to environmental hazards while having limited access to resources that could help them cope with or avoid these exposures (Malin & Ryder, 2018). These disparities are often compounded by

climate change, which tends to exacerbate existing inequalities and create new forms of environmental injustice (Islam & Winkel, 2017). Conceptual frameworks that fail to account for these equity dimensions provide an incomplete picture of environmental health impacts and may inadvertently perpetuate health disparities (Schlosberg & Collins, 2014).

The complexity of environmental health challenges has driven the development of systems-based approaches that recognize the interconnected nature of environmental, social, and health systems (McMichael, 2013). These approaches emphasize the importance of understanding feedback loops, tipping points, and emergent properties that arise from the interaction of multiple system components (Bai *et al.*, 2016). Systems thinking has proven particularly valuable in addressing climate change and health linkages, where the impacts of environmental changes cascade through multiple pathways and scales to influence health outcomes (Hosking & Campbell-Lendrum, 2012).

The role of technology and data analytics in environmental health has expanded dramatically, creating new opportunities exposure assessment, health surveillance, and intervention design (Nwankwo et al., 2024). Advanced modeling techniques, remote sensing technologies, and big data approaches are enabling more sophisticated analyses of environment-health relationships while supporting the development of predictive models that can inform proactive public health responses (Akinboboye et al., 2022). These technological advances are particularly important for addressing climate-related health risks, where early warning systems and adaptive management approaches are essential for effective prevention strategies (Frempong et al., 2022). Public health practice is increasingly recognizing the importance of place-based approaches that account for the unique environmental, social, and economic characteristics of specific communities (O'Neill & Abson, 2009). These approaches acknowledge that environmental health interventions must be tailored to local contexts and that community engagement is essential for developing effective and sustainable solutions (Israel et al., 2012). Place-based frameworks also recognize the importance of local knowledge and community capacity in identifying environmental health risks and developing appropriate response strategies (Corburn, 2005).

The integration of environmental health considerations into broader health promotion and disease prevention strategies represents an important evolution in public health practice (Hancock, 2015). Rather than treating environmental factors as separate from other health determinants, contemporary approaches recognize the need for comprehensive strategies that address multiple determinants simultaneously (Marmot & Wilkinson, 2006). This integrated approach is particularly important for addressing climate change and health linkages, where effective responses require coordination across multiple sectors and scales of action (Woodcock *et al.*, 2009).

#### 2. Literature Review

The literature on environmental health and disease prevention has expanded exponentially over the past two decades, reflecting growing recognition of the critical role that environmental factors play in shaping population health outcomes (Landrigan *et al.*, 2018). Early environmental health research focused primarily on occupational exposures and acute effects of high-level pollution exposures, but the

field has evolved to encompass a much broader range of environmental factors and health outcomes (Brunekreef & Holgate, 2002). Contemporary research addresses complex exposure scenarios involving multiple pollutants, chronic low-level exposures, and the interaction of environmental factors with genetic, behavioral, and social determinants of health (Wild, 2005).

The conceptual frameworks that guide environmental health research have undergone significant refinement as the field has matured (Okoye *et al.*, 2024). Early models were largely based on toxicological paradigms that emphasized doseresponse relationships and threshold effects, but these have been supplemented by more sophisticated frameworks that account for population heterogeneity, cumulative exposures, and the role of social determinants in modifying environmental health impacts (Sexton & Hattis, 2007). The development of the exposome concept represents a particularly important advancement, providing a framework for considering the totality of environmental exposures across the lifespan and their interactions with internal biological processes (Wild, 2012).

Air pollution research has provided some of the most compelling evidence for the health impacts of environmental exposures, with studies demonstrating associations between ambient air pollution and a wide range of health outcomes including cardiovascular disease, respiratory disease, cancer, and premature mortality (Pope et al., 2020). The Global Burden of Disease project has identified air pollution as one of the leading risk factors for disease burden globally, with particular impacts on vulnerable populations including children, elderly adults, and individuals with pre-existing health conditions (Cohen et al., 2017). Research has increasingly focused on understanding the mechanisms through which air pollution affects health, with evidence pointing to systemic inflammation, oxidative stress, and epigenetic modifications as key pathways (Rajagopalan et al., 2018).

Water quality and sanitation represent fundamental environmental health determinants, with substantial evidence linking inadequate water and sanitation services to infectious disease transmission, malnutrition, and impaired child development (Prüss-Ustün et al., 2019). The literature demonstrates that water-related health impacts extend beyond infectious diseases to include chemical contamination effects, with particular concerns about emerging contaminants such as pharmaceuticals, personal products, and endocrine-disrupting chemicals (Richardson & Ternes, 2018). Climate change is altering water availability and quality patterns, creating new challenges for water security and associated health outcomes (Howard et al., 2016).

Chemical exposure research has expanded to address the health impacts of thousands of synthetic chemicals that are now present in the environment, with particular attention to persistent organic pollutants, heavy metals, and endocrine-disrupting chemicals (Grandjean & Bellanger, 2017). The literature reveals that chemical exposures often occur as complex mixtures, making it difficult to isolate the effects of individual chemicals and necessitating new approaches to mixture risk assessment (Kortenkamp *et al.*, 2009). Research on developmental origins of health and disease has highlighted the particular vulnerability of early life stages to chemical exposures, with evidence that exposures during critical developmental windows can have long-lasting effects

on health outcomes (Heindel et al., 2017).

The climate change and health literature has grown rapidly, with comprehensive assessments identifying multiple pathways through which climate change affects human health (Watts et al., 2021). Direct health effects include temperature-related morbidity and mortality, with evidence showing increased risks of heat-related cardiovascular events, and respiratory complications during extreme heat events (Basu, 2009). Indirect health effects operate through multiple pathways including altered patterns of infectious disease transmission, food security impacts, extreme weather events, and ecosystem disruption (McMichael et al., 2012). The literature increasingly recognizes that climate change acts as a threat multiplier, exacerbating existing environmental health risks while creating new exposure scenarios (Hayes et al., 2018).

Environmental justice research has documented substantial disparities in environmental health impacts across racial, ethnic, and socioeconomic groups, with marginalized communities often facing disproportionate exposure to environmental hazards (Malin & Ryder, 2018). The literature reveals that these disparities arise through multiple mechanisms including discriminatory land use decisions, differential access to resources for avoiding or coping with exposures, and cumulative impacts of multiple stressors (Morello-Frosch & Lopez, 2006). Climate change is expected to exacerbate these existing disparities, with vulnerable populations facing greater risks from climate-related health impacts while having limited capacity for adaptation (Islam & Winkel, 2017).

Systems approaches to environmental health have gained prominence in the literature, with researchers recognizing the need for frameworks that can capture the complex interactions between environmental, social, and health systems (McMichael, 2013). These approaches emphasize the importance of understanding feedback loops, non-linear relationships, and emergent properties that arise from system interactions (Bai *et al.*, 2016). The literature on social-ecological systems has provided valuable insights into how human and natural systems interact to influence health outcomes, with particular relevance for understanding climate change and health linkages (Folke *et al.*, 2010).

The role of urban environments in shaping health outcomes has received increasing attention in the literature, with research demonstrating that urban design and planning decisions have profound impacts on environmental exposures and health outcomes (Frumkin *et al.*, 2004). The concept of healthy cities has emerged as an important framework for integrating health considerations into urban planning processes, with evidence showing that well-designed urban environments can promote health while reducing environmental risks (Rydin *et al.*, 2012). Research on urban heat islands, air quality, and green space access has provided important insights into the mechanisms through which urban environments influence health (Akhamere, 2023).

The literature on environmental health interventions has evolved from a focus on end-of-pipe pollution control to more comprehensive approaches that address the root causes of environmental health problems (Friel *et al.*, 2011). Primary prevention strategies that eliminate or reduce environmental exposures at their source are increasingly recognized as more effective and cost-efficient than approaches that focus on treating health effects after exposure has occurred (Grandjean, 2013). The literature emphasizes the importance

of multi-sectoral collaboration, community engagement, and policy interventions that address the structural determinants of environmental health disparities (Corburn, 2005).

Technological advances in environmental monitoring, exposure assessment, and health surveillance have created new opportunities for advancing environmental health research and practice (Appoh *et al.*, 2022). The literature on environmental health informatics demonstrates how big data approaches, remote sensing technologies, and advanced modeling techniques can enhance our ability to understand and address environmental health challenges (Jerrett, 2009). These technological tools are particularly important for addressing climate change and health linkages, where real-time monitoring and predictive modeling are essential for effective adaptation strategies (Umana *et al.*, 2022).

The integration of environmental health considerations into broader public health and health care practice represents an important theme in recent literature (Hancock, 2015). Research has demonstrated that health care systems have important roles to play in addressing environmental health challenges, both through direct patient care and through advocacy and policy engagement (Sheffield & Landrigan, 2011). The literature on planetary health has emerged as an important framework for understanding the connections between human health and the health of natural systems, emphasizing the need for approaches that address both human and environmental wellbeing (Whitmee *et al.*, 2015).

#### 3. Methodology

This comprehensive review employed a systematic approach to examine the conceptual frameworks linking pollution exposure, climate change, and public health outcomes in the context of environmental health and disease prevention. The methodology was designed to provide a thorough analysis of current knowledge while identifying key gaps and opportunities for advancing the field. The research approach integrated multiple methodological strategies including systematic literature review, conceptual framework analysis, and evidence synthesis to develop a comprehensive understanding of the complex relationships between environmental factors and health outcomes (Okoli *et al.*, 2022).

The literature search strategy was developed using a combination of key terms related to environmental health, pollution exposure, climate change, and public health outcomes. Primary databases searched included PubMed, Web of Science, Scopus, and Environmental Index, with additional searches conducted in specialized databases such as TOXLINE and GreenFILE. The search strategy was designed to capture both empirical research studies and theoretical papers that contribute to our understanding of environment-health relationships. Search terms were combined using Boolean operators to create comprehensive search strings that could identify relevant literature across multiple domains of environmental health research (Afrihyia et al. 2022)

Inclusion criteria for the literature review were established to focus on peer-reviewed articles published between 2000 and 2024 that addressed environmental health determinants, pollution exposures, climate change impacts, or public health outcomes related to environmental factors. Studies were included if they provided empirical evidence of environmenthealth relationships, presented conceptual frameworks for understanding these relationships, or offered methodological

advances relevant to environmental health research. Both quantitative and qualitative studies were included to ensure comprehensive coverage of the literature. Priority was given to studies that addressed multiple environmental exposures, vulnerable populations, or intervention strategies (Omolayo *et al.*, 2023).

The conceptual framework analysis involved systematic examination of the theoretical models and frameworks used in environmental health research to understand environmenthealth relationships. This analysis focused on identifying common elements across frameworks, assessing the strengths and limitations of different approaches, and examining how frameworks have evolved over time to address emerging challenges such as climate change and environmental justice. The analysis also examined how different frameworks address issues of scale, complexity, and uncertainty that are inherent in environmental health research (Frempong *et al.*, 2024).

Data extraction was conducted using standardized forms that captured key information about study design, population characteristics, environmental exposures, health outcomes, and study findings. For conceptual framework analysis, data extraction focused on identifying the key components of each framework, the relationships between components, and the underlying assumptions and theories that inform the framework. Quality assessment was conducted using established criteria appropriate for different study designs, with particular attention to issues of exposure assessment quality, confounding control, and generalizability of findings (Akhamere, 2022).

The evidence synthesis approach involved both narrative synthesis and where appropriate, quantitative synthesis of findings across studies. The narrative synthesis focused on identifying patterns in the evidence, assessing the consistency of findings across different contexts and populations, and examining factors that might explain heterogeneity in results. The synthesis was structured around key themes including direct health effects of environmental exposures, indirect effects operating through ecosystem disruption, the role of social determinants in modifying environmental health impacts, and the implications of climate change for environmental health risks (Dogho, 2023).

Geographic and temporal considerations were integrated into the methodology to ensure that the analysis captured variation in environmental health relationships across different contexts and time periods. The review included studies from diverse geographic regions including both developed and developing countries, with particular attention to studies from Africa and other regions that may face unique environmental health challenges. Temporal analysis examined how environmental health risks and conceptual frameworks have evolved over time, with special focus on how climate change has altered our understanding of environment-health relationships (Ayumu & Ohakawa, 2024).

Population vulnerability analysis was incorporated as a key methodological component, with systematic examination of how environmental health impacts vary across different demographic groups including children, elderly adults, pregnant women, and socioeconomically disadvantaged populations. This analysis examined both biological factors that may increase susceptibility to environmental exposures and social factors that may increase exposure risk or reduce capacity to cope with environmental health threats. The

analysis also examined how climate change may alter patterns of vulnerability and create new at-risk populations (Olajide *et al.*, 2023).

The methodology included systematic assessment of intervention strategies and their effectiveness in addressing environmental health challenges. This analysis examined both traditional pollution control approaches and more innovative strategies that address the social and structural determinants of environmental health disparities. The assessment considered the scalability, sustainability, and equity implications of different intervention approaches, with particular attention to strategies that could be effective in addressing climate-related health risks (Alonge et al., 2023). Quality assurance procedures were implemented throughout the methodology to ensure the reliability and validity of the findings. These procedures included independent screening and data extraction by multiple reviewers, regular calibration exercises to ensure consistency in data extraction and quality assessment, and systematic documentation of methodological decisions and potential sources of bias. The analysis also included sensitivity analyses to examine the robustness of findings to different methodological choices and assumptions (Ilori, 2022).

The methodology incorporated stakeholder engagement strategies to ensure that the analysis addressed the needs and priorities of environmental health practitioners, policymakers, and affected communities. This engagement involved consultation with experts in environmental health, climate change, and public health to validate the analytical framework and ensure that the synthesis addressed key policy and practice questions. Community stakeholder input was also incorporated to ensure that the analysis reflected the perspectives and experiences of populations most affected by environmental health challenges (Fagbore *et al.*, 2024).

### **3.1.** Theoretical Foundations of Environmental Health Frameworks

The theoretical foundations of environmental health frameworks have evolved substantially over the past several decades, reflecting advances in our understanding of exposure pathways, biological mechanisms, and the complex interactions between environmental, social, and individual factors that determine health outcomes. Early environmental health frameworks were largely grounded in toxicological models that emphasized linear dose-response relationships and assumed that health effects could be predicted based on exposure levels and toxicity data (Sexton & Hattis, 2007). These models provided important insights into acute health effects of high-level exposures but proved inadequate for understanding the complex, multi-factorial nature of environmental health impacts in real-world settings where people are exposed to multiple pollutants at varying levels over extended periods.

The development of epidemiological approaches to environmental health research introduced new theoretical perspectives that emphasized population-level patterns of disease and the role of environmental factors as determinants of health across populations (Rothman *et al.*, 2008). Epidemiological frameworks brought attention to issues of confounding, effect modification, and the challenges of establishing causal relationships between environmental exposures and health outcomes in observational studies. These frameworks also highlighted the importance of vulnerable populations and individual susceptibility factors

that could modify the relationship between exposure and health outcomes (Vrijheid, 2014). The integration of epidemiological and toxicological perspectives has been essential for developing comprehensive frameworks that can address both mechanistic understanding and population health impacts.

Systems thinking has emerged as a fundamental theoretical foundation for contemporary environmental health frameworks, recognizing that environmental health outcomes arise from complex interactions within and between multiple systems including environmental, social, economic, and biological systems (McMichael, 2013). Systems approaches emphasize the importance of feedback loops, non-linear relationships, and emergent properties that cannot be understood by examining individual system components in isolation. This perspective has been particularly valuable for understanding climate change and health linkages, where environmental changes cascade through multiple pathways to influence health outcomes at different temporal and spatial scales (Bai *et al.*, 2016).

The concept of environmental justice has provided crucial theoretical grounding for frameworks that address the social determinants of environmental health disparities (Bullard & Johnson, 2000). Environmental justice theories emphasize that environmental health impacts are not randomly distributed across populations but are shaped by social, economic, and political factors that influence both exposure patterns and vulnerability to environmental health risks. These theories highlight the importance of procedural justice in environmental decision-making, distributive justice in the allocation of environmental benefits and burdens, and recognition justice in acknowledging the knowledge and experiences of affected communities (Schlosberg, 2007).


The exposome concept represents a significant theoretical advancement that provides a framework for considering the totality of environmental exposures across the lifespan and their interactions with internal biological processes (Wild, 2012). This concept recognizes that individuals are exposed to complex mixtures of environmental agents that may interact with each other and with individual biological characteristics to influence health outcomes. The exposome framework emphasizes the need for comprehensive exposure assessment approaches that can capture both external environmental exposures and internal biological responses to those exposures. This perspective has important implications for understanding cumulative health impacts and developing personalized approaches to environmental health protection (Vermeulen *et al.*, 2020).

Life course epidemiology has provided theoretical foundations for understanding how environmental exposures at different life stages contribute to health outcomes across the lifespan (Ben-Shlomo & Kuh, 2002). This theoretical perspective emphasizes that certain developmental periods may be particularly sensitive to environmental exposures, with effects that may not manifest until later in life. The developmental origins of health and disease hypothesis has been particularly influential, providing evidence that environmental exposures during critical developmental windows can have long-lasting effects on health outcomes (Heindel *et al.*, 2017). These concepts have important implications for environmental health policy and intervention strategies, emphasizing the need for protection of vulnerable life stages.

Social-ecological systems theory has provided important

theoretical foundations for understanding the interactions between human and natural systems that influence environmental health outcomes (Folke *et al.*, 2010). This theoretical perspective recognizes that human health is fundamentally dependent on the health of natural systems and that environmental health challenges cannot be understood without considering the social, economic, and political factors that influence human-environment interactions. Social-ecological systems theory emphasizes concepts such as resilience, adaptive capacity, and transformation that are essential for understanding how communities can respond to environmental health challenges, particularly in the context of climate change (Norberg & Cumming, 2008).

The precautionary principle has emerged as an important theoretical foundation for environmental health policy and practice, particularly in situations where scientific uncertainty exists about the health effects of environmental exposures (Kriebel *et al.*, 2001). This principle suggests that protective action should be taken even in the absence of complete scientific certainty when potential health impacts are serious or irreversible. The precautionary principle has been particularly important for addressing emerging environmental health threats such as endocrine-disrupting chemicals and nanoparticles, where traditional risk assessment approaches may be inadequate for protecting public health (Grandjean, 2013).



**Fig 1:** Theoretical Evolution of Environmental Health Frameworks (2000-2024)

Risk assessment frameworks have provided important theoretical foundations for evaluating environmental health hazards and informing regulatory decision-making (NRC, 2009). Traditional risk assessment approaches involve hazard identification, dose-response assessment, exposure assessment, and risk characterization, providing a systematic

approach for evaluating the potential health impacts of environmental exposures. However, these frameworks have been critiqued for their focus on single chemicals, their reliance on animal toxicity data, and their limited consideration of vulnerable populations and cumulative exposures (Sexton *et al.*, 2004). Contemporary risk assessment frameworks are evolving to address these limitations through approaches such as cumulative risk assessment, aggregate exposure assessment, and community-based participatory risk assessment.

Health impact assessment has emerged as an important theoretical framework for evaluating the potential health consequences of policies, programs, and projects that may affect environmental conditions (Dannenberg *et al.*, 2008). This framework provides a systematic approach for identifying, predicting, and evaluating the health impacts of proposed actions, with particular attention to impacts on vulnerable populations and health equity considerations. Health impact assessment frameworks emphasize stakeholder engagement, interdisciplinary collaboration, and the use of multiple types of evidence including quantitative data, qualitative information, and community knowledge (Harris-Roxas *et al.*, 2012).

One Health approaches have provided theoretical foundations for understanding the interconnections between human health, animal health, and environmental health (Zinsstag *et al.*, 2011). This perspective recognizes that human health cannot be understood in isolation from the health of animals and ecosystems and emphasizes the need for collaborative approaches that address health challenges across these domains. One Health frameworks have been particularly valuable for understanding zoonotic disease transmission, antimicrobial resistance, and food safety issues that arise at the interface of human, animal, and environmental health systems (Ruckelshaus *et al.*, 2020).

### 3.2. Pollution Exposure Pathways and Health Impact Mechanisms

Environmental pollution exposures operate through complex pathways that involve multiple environmental media, exposure routes, and biological mechanisms that ultimately influence health outcomes across diverse populations. Understanding these exposure pathways is fundamental to developing effective environmental health frameworks and designing appropriate intervention strategies. The complexity of real-world exposures, which typically involve simultaneous exposure to multiple pollutants through multiple routes over varying time periods, requires sophisticated analytical approaches that can capture the cumulative and interactive effects of these exposures (Sexton & Hattis, 2007).

Air pollution represents one of the most significant environmental health threats globally, with exposure pathways involving inhalation of particulate matter, gaseous pollutants, and complex mixtures of atmospheric contaminants (Pope *et al.*, 2020). Fine particulate matter (PM2.5) has received particular attention due to its ability to penetrate deep into the respiratory system and enter the bloodstream, where it can trigger systemic inflammatory responses and contribute to cardiovascular and respiratory disease (Rajagopalan *et al.*, 2018). The mechanisms through which air pollution affects health include oxidative stress, systemic inflammation, endothelial dysfunction, and epigenetic modifications that can alter gene expression

patterns and contribute to disease development (Brook *et al.*, 2010). These mechanisms help explain why air pollution exposure is associated with such a wide range of health outcomes, from acute respiratory symptoms to chronic diseases and premature mortality.

Water contamination pathways involve exposure to chemical, biological, and physical contaminants through drinking water consumption, recreational water contact, and food preparation activities (Richardson & Ternes, 2018). Chemical contaminants in water supplies include heavy metals, organic pollutants, disinfection byproducts, and emerging contaminants such as pharmaceuticals and personal care products. The health impacts of water contamination operate through multiple mechanisms including direct toxicity, disruption of endocrine function, carcinogenic effects, and increased susceptibility to infectious diseases (Prüss-Ustün *et al.*, 2019). Biological contaminants in water systems can cause acute gastroenteritis and other infectious diseases, with particular risks for vulnerable populations including children, elderly adults, and immunocompromised individuals

Soil contamination represents another important exposure pathway, with human exposure occurring through direct contact with contaminated soil, inhalation of dust particles, and consumption of contaminated food grown in polluted soil (Swartjes, 2011). Heavy metals such as lead, cadmium, and arsenic can accumulate in soil from industrial activities, mining operations, and agricultural practices, creating long-term exposure risks for nearby communities. The health impacts of soil contamination are particularly concerning for children, who may have higher exposure rates due to hand-to-mouth behavior and may be more susceptible to the developmental effects of toxic exposures (Lanphear *et al.*, 2016). The persistence of many soil contaminants means that exposure risks can continue for decades or centuries after the original contamination occurred.

Food chain contamination pathways involve the accumulation and concentration of pollutants as they move

through the food web, with particular concerns about persistent organic pollutants and heavy metals that bioaccumulate in fatty tissues (Schecter *et al.*, 2010). Seafood consumption represents a major exposure pathway for mercury, which can cause neurological damage, particularly during fetal development. Agricultural practices can introduce pesticides, antibiotics, and other contaminants into the food supply, creating widespread population exposures to these chemicals. The globalization of food systems has increased the complexity of food chain contamination, as pollutants released in one region can affect food supplies consumed in distant locations (Nwankwo *et al.*, 2024).

Indoor air pollution represents a significant exposure pathway that is often overlooked in environmental health assessments, despite the fact that people spend the majority of their time indoors (Jones, 1999). Indoor pollutants include combustion products from cooking and heating, volatile organic compounds from building materials and consumer products, biological contaminants such as mold and dust mites, and outdoor pollutants that infiltrate into indoor spaces. The health impacts of indoor air pollution can be substantial, particularly for vulnerable populations who spend more time indoors, such as young children, elderly adults, and individuals with chronic diseases (Mendell, 2007).

Occupational exposure pathways represent a critical component of environmental health, with workers often facing higher levels of exposure to toxic substances than the general population (Landrigan *et al.*, 2018). Occupational exposures can occur through inhalation, dermal contact, and ingestion of workplace contaminants, with health impacts ranging from acute poisoning to chronic diseases such as cancer, respiratory disease, and neurological disorders. The health effects of occupational exposures extend beyond individual workers to affect families and communities through take-home exposures and environmental releases from industrial facilities (Grandjean & Bellanger, 2017).

Table 1: Major Environmental Pollution Exposure Pathways and Associated Health Mechanisms

Exposure Pathway	Primary Contaminants	<b>Exposure Route</b>	Key Health Mechanisms	Vulnerable Populations
Air Pollution	PM2.5, NO2, O3, VOCs	Inhalation	Oxidative stress, systemic inflammation, endothelial dysfunction	Children, elderly, respiratory/cardiovascular disease patients
Water Contamination	Heavy metals, pathogens, disinfection byproducts, pharmaceuticals	Ingestion, dermal contact	Direct toxicity, endocrine disruption, infectious disease	Children, pregnant women, immunocompromised individuals
Soil Contamination	Lead, cadmium, arsenic, petroleum products	Ingestion, dermal contact, inhalation	Neurotoxicity, carcinogenesis, developmental effects	Children, agricultural workers, urban residents
Food Chain	Mercury, pesticides, persistent organic pollutants	Ingestion	Bioaccumulation, neurotoxicity, endocrine disruption	Pregnant women, high seafood consumers, subsistence communities
Indoor Air	VOCs, combustion products, biological agents	Inhalation	Respiratory irritation, sensitization, carcinogenesis	Households using solid fuels, children, asthmatic individuals
Occupational	Industrial chemicals, dusts, metals	Inhalation, dermal, ingestion	Acute toxicity, chronic disease, carcinogenesis	Industrial workers, agricultural workers, healthcare workers

Environmental mixture effects represent a particularly important consideration in understanding pollution exposure pathways, as real-world exposures typically involve simultaneous exposure to multiple pollutants that may interact in complex ways (Kortenkamp *et al.*, 2009). These interactions can result in additive effects, where the combined effect equals the sum of individual effects, synergistic effects where the combined effect is greater than the sum of

individual effects, or antagonistic effects where one pollutant reduces the toxicity of another. Understanding mixture effects is crucial for accurate risk assessment and for developing effective intervention strategies that address the full complexity of environmental exposures (Rider *et al.*, 2018).

The temporal dimension of exposure pathways is critical for understanding health impacts, as the timing, duration, and frequency of exposures can significantly influence the magnitude and nature of health effects (Ben-Shlomo & Kuh, 2002). Acute exposures may cause immediate health effects such as respiratory irritation or poisoning, while chronic exposures may contribute to the development of cancer, cardiovascular disease, or other chronic conditions over many years or decades. Critical windows of exposure, particularly during fetal development and early childhood, may have particularly significant impacts on health outcomes that may not manifest until later in life (Heindel et al., 2017). Geographic variation in exposure pathways reflects differences in pollution sources, environmental conditions, and population characteristics that influence both exposure levels and health impacts (Jerrett, 2009). Urban areas may have higher levels of air pollution from traffic and industrial sources, while rural areas may have greater exposures to agricultural chemicals and naturally occurring contaminants. Developing countries may face unique exposure scenarios related to industrial development, inadequate environmental controls, and limited access to clean water and sanitation (Landrigan et al., 2018). Understanding these geographic patterns is essential for developing targeted intervention strategies and addressing environmental health disparities. The role of individual susceptibility factors in modifying exposure-health relationships represents an important area of

The role of individual susceptibility factors in modifying exposure-health relationships represents an important area of environmental health research, with evidence showing that genetic factors, nutritional status, pre-existing health conditions, and other individual characteristics can influence how people respond to environmental exposures (Vrijheid, 2014). These susceptibility factors help explain why some individuals may experience health effects from environmental exposures while others do not, and why certain populations may be more vulnerable to environmental health risks. Understanding individual susceptibility is important for developing personalized approaches to environmental health protection and for identifying populations that may need additional protection from environmental exposures.

### 3.3. Climate Change as an Environmental Health Determinant

Climate change has emerged as one of the most significant environmental health challenges of the 21st century, fundamentally altering the environmental conditions that determine human health outcomes and creating new pathways through which environmental factors influence disease patterns (Watts *et al.*, 2021). The health impacts of climate change operate through multiple interconnected mechanisms that span direct physiological effects, indirect effects mediated through environmental and social systems, and complex feedback loops that can amplify health risks over time. Understanding these climate-health linkages requires conceptual frameworks that can capture the dynamic, multi-scale, and often non-linear nature of climate impacts on human health (McMichael *et al.*, 2012).

Direct health effects of climate change include temperaturerelated morbidity and mortality, which have become increasingly evident as global temperatures continue to rise and extreme heat events become more frequent and intense (Basu, 2009). Heat-related health impacts operate through multiple physiological pathways including heat stress, dehydration, exacerbation of cardiovascular and respiratory conditions, and heat stroke in severe cases. Vulnerable populations including elderly adults, young children, individuals with chronic diseases, and outdoor workers face disproportionate risks from extreme heat exposure. Urban heat island effects can amplify these risks in densely populated areas, where built environments retain heat and create temperature differentials that can exceed surrounding rural areas by several degrees (Rizwan *et al.*, 2008).

Extreme weather events represent another direct pathway through which climate change affects health, with increasing frequency and intensity of hurricanes, floods, droughts, and other extreme events creating immediate and long-term health risks (Hayes *et al.*, 2018). The health impacts of extreme weather events include injuries and deaths from the events themselves, mental health impacts from trauma and displacement, infectious disease outbreaks related to disrupted water and sanitation systems, and chronic health effects from exposure to contaminated flood waters or other environmental hazards. The recovery period following extreme weather events can be prolonged, with health impacts persisting for months or years after the initial event (Goldmann & Galea, 2014).

Air quality impacts of climate change create important indirect pathways for health effects, as rising temperatures and changing weather patterns alter the formation and distribution of air pollutants (Jacob & Winner, 2009). Climate change can increase ground-level ozone formation through temperature-dependent photochemical reactions, extend pollen seasons and increase allergen exposure, and alter the transport and dispersion of particulate matter and other air pollutants. Wildfires, which are becoming more frequent and severe due to climate change, represent a particularly important source of air pollution that can affect air quality over large geographic areas and create health risks for millions of people (Reid *et al.*, 2016).

Water security and quality impacts of climate change affect health through multiple pathways including altered precipitation patterns, increased flooding and drought, rising sea levels, and temperature-related changes in water quality (Howard *et al.*, 2016). Drought conditions can concentrate pollutants in water supplies and reduce access to safe drinking water, while flooding can contaminate water supplies with pathogens, chemicals, and other hazardous substances. Rising sea levels can lead to saltwater intrusion into freshwater supplies, making them unsuitable for drinking and agriculture. Changes in water temperature can alter the growth of waterborne pathogens and create new risks for waterborne disease outbreaks (Hunter, 2003).

Food security and nutrition impacts represent critical pathways through which climate change affects health, as changing temperature and precipitation patterns alter agricultural productivity and food availability (Wheeler & von Braun, 2013). Heat stress, drought, and extreme weather events can reduce crop yields and livestock productivity, leading to food shortages and increased food prices that disproportionately affect low-income populations. Climate change can also alter the nutritional quality of food crops, with elevated carbon dioxide concentrations reducing protein and micronutrient content in many staple crops (Myers *et al.*, 2014). These changes in food security and nutrition can have cascading effects on child development, maternal health, and population health outcomes.

Vector-borne disease transmission represents an important climate-sensitive health outcome, as changing temperature and precipitation patterns alter the geographic distribution, seasonal patterns, and transmission intensity of diseases such as malaria, dengue fever, and Lyme disease (Ogden & Lindsay, 2016). Warmer temperatures can expand the geographic range of disease vectors such as mosquitoes and ticks, while changes in precipitation can create new breeding habitats or eliminate existing ones. The relationships between climate and vector-borne disease transmission are complex and vary by disease system, but there is growing evidence that climate change is already altering disease transmission patterns in many regions (Semenza & Menne, 2009).

Mental health impacts of climate change operate through multiple pathways including direct effects of extreme weather events, indirect effects of environmental degradation and displacement, and broader psychological impacts related to climate anxiety and concern about future climate impacts (Clayton, 2020). Natural disasters can cause acute trauma and lead to post-traumatic stress disorder, depression, and other mental health conditions. Slow-onset climate impacts such as drought, sea level rise, and ecosystem degradation can cause chronic stress and contribute to mental health problems, particularly in communities whose livelihoods depend on climate-sensitive resources. Climate change can also exacerbate existing mental health conditions and create new stressors that affect psychological wellbeing (Palinkas, 2020).




Fig 2: Multi-Pathway Framework for Climate Change Health Impacts

Ecosystem health impacts of climate change affect human health through the disruption of ecosystem services that support human wellbeing, including clean air and water provision, climate regulation, pollination services, and (Millennium hazard protection Ecosystem Assessment, 2005). Climate change can alter ecosystem structure and function, leading to biodiversity loss, habitat degradation, and reduced capacity of natural systems to provide these essential services. Coastal ecosystems such as coral reefs and mangroves provide important protection from storm surge and flooding, and their degradation due to climate change can increase vulnerability to extreme weather events (Barbier et al., 2008).

Social and economic disruption caused by climate change creates additional pathways for health impacts, as climate-related damages to infrastructure, agriculture, and other economic sectors can affect employment, income, and access to health care and other essential services (Berry *et al.*, 2018). Climate-related migration and displacement can disrupt social networks and community support systems that are important for health and wellbeing. These social and economic impacts often disproportionately affect vulnerable populations who have limited resources to adapt to changing conditions (Islam & Winkel, 2017).

The temporal and spatial scales at which climate change operates present unique challenges for environmental health assessment and response, as climate impacts can manifest across multiple time scales from immediate effects during extreme events to gradual changes that unfold over decades or centuries (Friel *et al.*, 2011). Similarly, climate impacts can operate at scales ranging from local microclimates to global atmospheric systems, requiring analytical frameworks that can address cross-scale interactions and emergent properties that arise from complex system dynamics (O'Neill *et al.*, 2017).

Adaptation and mitigation strategies represent important components of climate change and health frameworks, as reducing greenhouse gas emissions and building adaptive capacity can significantly reduce health risks from climate change (Watts *et al.*, 2015). Health co-benefits of climate mitigation strategies, such as reduced air pollution from clean energy transitions and increased physical activity from active transportation, can provide immediate health benefits while also addressing long-term climate risks. Adaptation strategies that focus on building resilient health systems, improving early warning systems, and strengthening community capacity can help reduce vulnerability to climate-related health risks (Ebi *et al.*, 2018).

### 3.4. Social Determinants and Environmental Health Equity

The intersection of social determinants and environmental health represents a critical area of inquiry that recognizes environmental health impacts are not distributed equally across populations, but are shaped by complex interactions between environmental exposures and social, economic, and political factors that influence both exposure patterns and vulnerability to environmental health risks (Malin & Ryder, 2018). Understanding these relationships is essential for developing effective environmental health interventions that can address the root causes of health disparities and promote environmental justice. The conceptual frameworks that guide this work must account for the structural factors that create and maintain environmental health inequities while also addressing the cumulative impacts of multiple stressors that disproportionately affect marginalized communities.

Environmental justice has emerged as a foundational concept for understanding how social determinants interact with environmental factors to create health disparities (Bullard & Johnson, 2000). Research has consistently demonstrated that communities of color, low-income populations, and other marginalized groups face disproportionate exposure to environmental hazards while having limited access to environmental amenities such as parks, clean air, and safe drinking water. These disparities arise through multiple

mechanisms including historical patterns of discriminatory land use planning, the siting of hazardous facilities in disadvantaged communities, and the differential enforcement of environmental regulations (Pellow, 2017). The cumulative effect of these disparities is that some communities face much higher environmental health risks than others, contributing to persistent health inequities across racial, ethnic, and socioeconomic lines.

Cumulative risk assessment has become an important framework for understanding how multiple environmental and social stressors interact to affect health outcomes in disadvantaged communities (Sexton & Hattis, 2007). Traditional environmental health approaches often focus on single exposures or single pollutants, but research has shown that many communities face simultaneous exposures to multiple environmental hazards combined with social stressors such as poverty, discrimination, and limited access to healthcare. These cumulative exposures can have additive or synergistic effects that result in health impacts that are greater than would be predicted from considering individual stressors in isolation. Cumulative risk frameworks recognize the need to assess the combined impact of chemical stressors, non-chemical stressors, and social vulnerability factors to accurately characterize health risks in disadvantaged communities (Morello-Frosch & Lopez, 2006).

Social vulnerability represents a key concept for understanding how social and economic factors influence susceptibility to environmental health risks (Cutter et al., 2003). Socially vulnerable populations may have higher exposure to environmental hazards due to factors such as residential location, occupational exposures, or limited ability to avoid contaminated areas. They may also be more susceptible to the health effects of environmental exposures due to factors such as poor nutritional status, limited access to healthcare, pre-existing health conditions, or psychosocial stress. Social vulnerability can also affect the ability to recover from environmental health impacts, with disadvantaged communities often facing longer recovery times and more severe long-term consequences from environmental disasters or contamination events (Flanagan et al., 2011).

The role of racism and discrimination in creating environmental health disparities has received increasing attention in recent years, with research demonstrating that racial bias in decision-making processes contributes to the disproportionate placement of environmental hazards in communities of color (Pulido, 2000). These patterns reflect both intentional discrimination and structural racism embedded in institutions and policies that systematically disadvantage racial and ethnic minorities. The health impacts

of racism extend beyond direct environmental exposures to include the physiological effects of chronic stress from experiencing discrimination, which can increase susceptibility to environmental health risks and contribute to health disparities (Williams & Mohammed, 2009).

Housing conditions represent a critical social determinant that influences environmental health outcomes, as poor housing quality can create or exacerbate environmental exposures while limiting the ability to avoid external environmental hazards (Jacobs et al., 2009). Substandard housing may have problems such as lead paint, mold, pest infestations, inadequate ventilation, or structural defects that create health risks for residents. Low-income households may have limited housing choices and may be more likely to live in areas with environmental hazards such as busy roads, industrial facilities, or contaminated sites. Energy poverty, where households cannot afford adequate heating or cooling, can create additional health risks and limit the ability to protect against extreme weather events (Hernández, 2016). Educational attainment and health literacy represent important social determinants that influence environmental health outcomes through multiple pathways (Schillinger et al., 2006). Higher levels of education may be associated with greater awareness of environmental health risks, better ability to access and understand health information, and greater capacity to take protective actions. Educational attainment is also associated with higher income and better employment opportunities, which can provide more resources for avoiding environmental exposures and accessing healthcare services. Health literacy specifically affects the ability to understand environmental health information and make informed decisions about protective behaviors (Finn & O'Fallon,

Employment and occupational factors represent critical social determinants that influence environmental health through both direct occupational exposures and indirect effects of job characteristics on health and wellbeing (Quinn & Kriebel, 2006). Workers in certain industries such as agriculture, construction, and manufacturing may face higher exposures to toxic substances, while workers in low-wage jobs may have limited ability to refuse unsafe work or advocate for better working conditions. Employment status affects access to health insurance and healthcare services, while job insecurity and work-related stress can affect health outcomes and susceptibility to environmental exposures. The concept of environmental health disparities in the workplace recognizes that occupational environmental health risks are not equally distributed across all workers, with certain groups facing disproportionate risks (Peckham et al., 2017).

Social Determinant	Mechanism of Influence	Examples of Environmental Health Impacts	Affected Populations	Intervention Approaches
Income/Poverty	Residential location, housing quality, access to resources	Living near pollution sources, poor housing conditions, limited healthcare access	Low-income communities, racial minorities	Affordable housing, economic development, healthcare access
Race/Ethnicity	Institutional discrimination, residential segregation	Disproportionate facility siting, environmental racism	Communities of color, immigrant populations	Civil rights enforcement, community organizing, policy reform
Education	Health literacy, risk awareness, advocacy capacity	Limited knowledge of risks, reduced protective behaviors	Less educated populations, limited English proficiency	Health education, community engagement, accessible information
Housing	Indoor environmental quality, neighborhood characteristics	Lead exposure, mold, proximity to hazards	Renters, low-income households, minorities	Housing code enforcement, weatherization, tenant protections
Employment	Occupational exposures, job-related stress, healthcare access	Workplace chemical exposure, limited safety protections	Blue-collar workers, undocumented workers, women	Occupational safety regulation, worker organizing, healthcare benefits
Age	Biological vulnerability, social resources, mobility	Increased susceptibility in children and elderly	Children, elderly adults	Age-specific protections, caregiver support, accessible services

Table 2: Social Determinants of Environmental Health Disparities and Their Mechanisms

Access to healthcare and social services represents an important social determinant that affects environmental health outcomes by influencing the ability to prevent, identify, and treat environmentally-related health conditions (Institute of Medicine, 2003). Communities with limited healthcare access may have difficulty obtaining screening for environmental exposures, receiving treatment for environmentally-related illnesses, or accessing preventive services that could reduce vulnerability to environmental health risks. Social services such as nutrition assistance, housing assistance, and transportation services can also affect environmental health by addressing social vulnerabilities that increase susceptibility to environmental exposures (Kawachi et al., 2002).

Community social capital and collective efficacy represent important social resources that can influence environmental health outcomes by affecting the ability of communities to identify environmental health problems, advocate for solutions, and implement protective measures (Sampson *et al.*, 1997). Strong social networks and community organizations can provide information about environmental health risks, support collective action to address environmental problems, and help individuals access resources for protection and recovery. Research has shown that communities with higher levels of social capital may be more successful in preventing the siting of environmental hazards and in obtaining cleanup of contaminated sites (Agyeman *et al.*, 2016).

The concept of intersectionality has become increasingly important for understanding environmental health disparities, recognizing that individuals and communities may face multiple, intersecting forms of disadvantage that create unique patterns of environmental health risk (Collins & Bilge, 2016). For example, low-income women of color may face environmental health risks that are different from those faced by middle-class white women or low-income white men, due to the intersection of gender, race, and class-based disadvantages. Intersectional approaches to environmental health recognize the need to understand and address these complex, overlapping forms of disadvantage rather than treating each social determinant as independent (Gee & Payne-Sturges, 2004).

Policy and institutional factors represent upstream social

determinants that influence environmental health disparities through their effects on environmental conditions, exposure patterns, and community vulnerability (Frieden, 2010). Environmental regulations, zoning laws, transportation policies, and other governmental decisions can have profound effects on environmental health outcomes and health equity. The effectiveness of environmental health policies may vary across communities depending on factors such as enforcement capacity, community engagement, and political representation. Environmental health frameworks increasingly recognize the need to address these upstream policy determinants in addition to individual and community-level factors (Northridge *et al.*, 2003).

### **3.5.** Challenges and Barriers to Environmental Health Integration

The integration of environmental health considerations into broader public health practice and policy faces numerous challenges and barriers that limit the effectiveness of current approaches and create obstacles to achieving optimal environmental health outcomes (Frumkin, 2005). These challenges operate at multiple levels, from individual practitioner and organizational capacity limitations to systemic issues related to institutional structures, funding mechanisms, and political priorities. Understanding and addressing these barriers is essential for advancing environmental health integration and developing more effective responses to complex environmental health challenges (Sheffield & Landrigan, 2011).

Institutional fragmentation represents one of the most significant barriers to environmental health integration, as environmental health responsibilities are typically divided across multiple agencies and organizations with different mandates, priorities, and approaches (Lowe *et al.*, 2013). Public health agencies may focus on health outcomes without adequate attention to environmental determinants, while environmental agencies may prioritize pollution control without sufficient consideration of health impacts. This fragmentation can lead to gaps in coverage, duplicated efforts, and missed opportunities for synergistic interventions that could address both environmental and health goals simultaneously. The lack of coordination between agencies can also create confusion for communities and stakeholders

about which agency is responsible for addressing specific environmental health problems (Corburn, 2005).

Limited capacity and training in environmental health among public health professionals represents another significant barrier to integration (Rudd & Buttke, 2012). Many public health practitioners lack adequate training in environmental health principles, exposure assessment methods, and environmental health intervention strategies. This knowledge gap can limit the ability to identify environmental health problems, conduct appropriate risk assessments, and develop effective intervention strategies. The complexity of environmental health issues, which often require understanding of toxicology, environmental science, epidemiology, and social determinants, creates challenges for practitioners who may not have interdisciplinary training (Burke *et al.*, 2017).

Funding limitations and competing priorities create significant barriers to environmental health integration, as environmental health programs often compete with other public health priorities for limited resources (Lichtveld et al., 2009). Environmental health interventions may require substantial upfront investments with benefits that accrue over long time periods, making them less attractive to policymakers focused on short-term outcomes. The preventive nature of many environmental interventions means that their benefits may not be immediately visible, making it difficult to demonstrate return on investment to funders and policymakers. Additionally, environmental health problems often require sustained, longterm interventions rather than the short-term project funding that characterizes much public health financing (Shendell et al., 2002).

Data limitations and surveillance gaps represent critical barriers to environmental health integration, as effective interventions require high-quality data on environmental exposures, health outcomes, and the relationships between them (Thacker et al., 2006). Environmental exposure data is often limited in geographic and temporal coverage, making it difficult to characterize population exposures accurately. surveillance systems may not environmentally-related health conditions adequately, particularly chronic diseases with long latency periods or subtle health effects that may not be recognized as environmentally-related. The lack of standardized data collection methods and data sharing protocols can further limit the ability to conduct comprehensive environmental health assessments (Teutsch & Thacker, 1995).

Scientific uncertainty and knowledge gaps create challenges for environmental health decision-making, as policymakers and practitioners may be reluctant to take action in the absence of definitive scientific evidence (Grandjean, 2013). The complexity of environmental health relationships, with multiple exposures, multiple health outcomes, and numerous confounding factors, makes it difficult to establish clear causal relationships in many cases. Emerging environmental health threats such as endocrine-disrupting chemicals, nanoparticles, and climate change present particular challenges due to limited long-term health data and uncertainty about exposure levels and health effects. The lag time between environmental exposures and health effects can make it difficult to establish causality and can delay recognition of emerging health threats (Woodruff et al., 2011).

Political and economic interests can create significant

barriers to environmental health integration, particularly when environmental health interventions conflict with economic interests or established political priorities (Wing, 2005). Industries may resist environmental health regulations that could increase costs or limit profitability, while political leaders may be reluctant to support interventions that could be seen as harmful to economic development. The influence of special interests in policy-making processes can limit the adoption of evidence-based environmental health policies and create barriers to effective enforcement of environmental health regulations. Community environmental health concerns may be discounted or ignored when they conflict with powerful economic or political interests (Brown, 1992). Community engagement and participation challenges create barriers to effective environmental health integration, as successful interventions often require active community involvement and support (Israel et al., 2012). Environmental health problems may not be recognized as priorities by affected communities, particularly when they compete with more immediate concerns such as employment, housing, or crime. Cultural and linguistic barriers can limit the effectiveness of environmental health communication and education efforts, while mistrust of government agencies and institutions can reduce community willingness to participate in environmental health programs. The technical complexity of environmental health issues can make it difficult for community members to understand and engage with environmental health information and decision-making processes (Corburn, 2005).

Legal and regulatory barriers can limit the effectiveness of environmental health integration efforts, particularly when existing laws and regulations are inadequate to address current environmental health challenges (Rechtschaffen & Markell, 2003). Environmental health laws may be outdated, fragmented across multiple statutes, or may lack adequate enforcement mechanisms. The burden of proof required to establish environmental health hazards may be set too high, making it difficult to take protective action in cases where scientific uncertainty exists. Jurisdictional limitations may prevent agencies from addressing environmental health problems that cross geographic or sectoral boundaries. Additionally, legal requirements for cost-benefit analysis may not adequately account for environmental health benefits, particularly those that accrue to disadvantaged communities (Heinzerling, 1998).

Technological and methodological limitations create challenges for environmental health assessment and intervention, particularly in areas such as exposure assessment, health surveillance, and risk assessment (Jerrett, 2009). Current exposure assessment methods may not capture the full complexity of real-world exposures, including cumulative exposures, mixture effects, and temporal variability in exposures. Health surveillance systems may not have the sensitivity to detect subtle environmental health effects or the geographic resolution to identify localized environmental health problems. Risk assessment methods may not adequately account for vulnerable populations, cumulative risks, or emerging health threats. The rapid pace of technological change can create challenges for regulatory systems that may not be able to keep pace with new technologies and associated health risks (Davies, 2009).

Communication and risk perception challenges create barriers to environmental health integration by affecting public understanding and acceptance of environmental health risks and interventions (Sandman, 1989). Environmental health risks may be difficult to communicate effectively due to their technical complexity, uncertainty, and long-term nature. Public risk perceptions may not align with scientific risk assessments, leading to situations where high-risk problems receive insufficient attention while low-risk problems generate disproportionate concern. Cultural, social, and psychological factors can influence how communities perceive and respond to environmental health risks, affecting the effectiveness of risk communication and intervention strategies. The politicization of environmental issues can further complicate communication efforts and reduce trust in environmental health information (Kasperson et al., 1988). Workforce development challenges create long-term barriers to environmental health integration, as the field requires specialized knowledge and skills that may not be adequately developed through current education and training programs (Lichtveld et al., 2006). Environmental health competencies span multiple disciplines including environmental science, toxicology, epidemiology, risk assessment, and policy analysis, requiring educational programs that can provide interdisciplinary training. The aging of the environmental health workforce and limited career advancement opportunities may create recruitment and retention challenges. Professional development opportunities for current practitioners may be limited, making it difficult to keep pace with rapidly evolving knowledge and methods in environmental health (Beck & Frankel, 2004).

### **3.6.** Best Practices and Recommendations for Framework Implementation

The development and implementation of effective environmental health frameworks require systematic approaches that integrate scientific evidence, community engagement, policy development, and institutional capacity building to address the complex challenges at the intersection of environmental factors and public health outcomes (Frumkin, 2010). Best practices in framework implementation have emerged from successful environmental health initiatives across diverse contexts, providing valuable insights into the strategies and approaches that can enhance the effectiveness of environmental health interventions while promoting equity and sustainability (Corburn, 2005). These practices emphasize the importance of adaptive management, multi-sectoral collaboration, and community-centered approaches that can respond to local needs and priorities while addressing systemic barriers to environmental health improvement.

Systems-based approaches represent a fundamental best practice for environmental health framework implementation, recognizing that environmental health challenges arise from complex interactions between environmental, social, economic, and political systems that require comprehensive intervention strategies (McMichael, 2013). Effective systems approaches involve mapping the relationships between different system components, identifying leverage points where interventions can have maximum impact, and developing coordinated strategies that address multiple system levels simultaneously. This approach requires moving beyond single-issue interventions to address the root causes of environmental health problems while building system capacity for long-term improvement. Systems approaches also emphasize the importance of monitoring and evaluation systems that can track progress across multiple outcomes and adapt strategies based on emerging evidence and changing conditions (Bai *et al.*, 2016).

Community-based participatory approaches have proven highly effective for environmental health framework implementation, particularly in addressing environmental justice concerns and ensuring that interventions are responsive to community needs and priorities (Israel et al., 2012). These approaches involve meaningful community engagement throughout all phases of environmental health work, from problem identification and research design to intervention implementation and evaluation. Communitybased approaches recognize that communities have valuable knowledge about local environmental conditions and health concerns that may not be captured through traditional scientific methods. They also emphasize capacity building within communities to enable ongoing environmental health advocacy and action. Successful community-based approaches require long-term commitments, flexible funding mechanisms, and institutional changes that can support authentic partnerships between communities and technical experts (Minkler & Wallerstein, 2012).

Integrated surveillance and monitoring systems represent critical infrastructure for effective environmental health framework implementation, providing the data needed to identify environmental health problems, track trends over time, and evaluate intervention effectiveness (Thacker et al., 2006). Best practices in surveillance system development emphasize the integration of environmental and health data, the use of multiple data sources to provide comprehensive assessments, and the development of real-time monitoring capabilities that can support rapid response to emerging threats. Modern surveillance systems increasingly utilize advanced technologies such as remote sensing, geographic information systems, and big data analytics to enhance data collection and analysis capabilities. Effective surveillance systems also require standardized data collection protocols, data sharing agreements, and quality assurance procedures to ensure data reliability and comparability across different contexts (Frempong et al., 2022).

Policy integration strategies represent essential components effective environmental health framework implementation, sustainable improvements as environmental health outcomes typically require supportive policy environments that can address systemic barriers and create incentives for protective actions (Frieden, 2010). Best practices in policy integration involve developing policy frameworks that address environmental health across sectors including health, transportation, housing, and economic development. This approach recognizes that many policy decisions outside the traditional health sector have important implications for environmental health outcomes. Effective policy integration also requires mechanisms for assessing the health impacts of proposed policies, ensuring that health considerations are incorporated into decision-making processes across sectors. Health in All Policies approaches provide valuable frameworks for achieving this integration (Rudolph et al.,

Capacity building and workforce development represent fundamental requirements for effective environmental health framework implementation, as successful interventions require skilled professionals who can address the technical, social, and political dimensions of environmental health challenges (Lichtveld et al., 2006). Best practices in capacity building emphasize interdisciplinary training that provides professionals with knowledge and skills across multiple domains including environmental science, public health, policy analysis, and community engagement. Professional development programs should provide ongoing opportunities for skill development and knowledge updating as the field evolves. Capacity building efforts should also focus on developing leadership skills and systems thinking capabilities that enable professionals to work effectively across organizational and sectoral boundaries. Community capacity building is equally important, providing community members with the knowledge and skills needed to participate effectively in environmental health decision-making and advocacy (Burke et al., 2017).

Evidence-based intervention strategies represent core components of effective environmental health framework implementation, requiring systematic approaches to identifying, evaluating, and scaling effective interventions (Brownson et al., 2009). Best practices in intervention development emphasize the importance of pilot testing and evaluation to assess effectiveness before large-scale implementation. Intervention strategies should be based on theoretical frameworks that explain how interventions are expected to achieve their intended outcomes, and should include process and outcome evaluation components that can provide feedback for continuous improvement. Effective interventions often involve multiple components that address different aspects of environmental health problems, requiring careful coordination and integration to achieve optimal outcomes. The scaling of successful interventions requires attention to contextual factors that may influence effectiveness in different settings (Glasgow et al., 2012).

Multi-sectoral collaboration and partnership development have emerged as critical success factors for environmental health framework implementation, as environmental health challenges typically require coordinated action across multiple organizations and sectors (Lowe et al., 2013). Effective partnerships involve clear agreements about roles, responsibilities, and expectations, as well as mechanisms for communication, coordination, and conflict resolution. Partnerships should include representation from all relevant stakeholders including government agencies, academic institutions, community organizations, and private sector entities. Long-term sustainability of partnerships requires ongoing attention to relationship building, shared decisionmaking processes, and equitable distribution of benefits and responsibilities. Successful partnerships often evolve over time, requiring adaptive management approaches that can respond to changing conditions and emerging opportunities (Lasker et al., 2001).

Technology integration and innovation represent increasingly important components of environmental health framework implementation, as new technologies create opportunities for more effective and efficient approaches to environmental health assessment and intervention (Jerrett, 2009). Best practices in technology integration emphasize the importance of selecting technologies that are appropriate for local contexts and user capabilities, while ensuring that technology solutions address real needs rather than creating new barriers. Mobile health technologies, environmental sensors, and data analytics platforms can enhance environmental health surveillance, education,

intervention delivery. However, technology integration must address issues of equity and accessibility to ensure that technological solutions do not exacerbate existing disparities. Privacy and data security considerations are also critical when implementing technology-based environmental health solutions (Akinboboye *et al.*, 2022).

Adaptive management approaches represent essential frameworks for environmental health implementation in contexts characterized by uncertainty, complexity, and changing conditions (Holling, 1978). Adaptive management involves systematic experimentation, monitoring, and learning to improve intervention effectiveness over time. This approach recognizes that environmental health interventions may need to be modified based on new evidence, changing conditions, or unforeseen consequences. Adaptive management requires organizational cultures that support learning and innovation, as well as management systems that can respond quickly to new information. Climate change adaptation represents a particularly important application of adaptive management approaches, as climate impacts are characterized by high uncertainty and changing risk profiles that require flexible response strategies (Ebi et al., 2016).

Financing and sustainability strategies represent critical considerations for environmental health framework implementation, as effective interventions often require sustained funding over long time periods to achieve meaningful health improvements (Shendell et al., 2002). Best practices in financing involve developing diversified funding strategies that combine public and private resources, while creating financing mechanisms that can support both shortterm interventions and long-term capacity building. Innovative financing approaches such as environmental health impact bonds, green bonds, and payment for ecosystem services can provide new resources for environmental health investments. Sustainability planning should be integrated into intervention design from the beginning, with attention to building local capacity, developing supportive policy environments, and creating economic incentives that support continued implementation. Cost-effectiveness analysis and economic evaluation can provide valuable information for demonstrating the value of environmental health investments to policymakers and funders (Ayumu & Ohakawa, 2024).

Quality assurance and performance measurement systems represent essential components of effective environmental health framework implementation, providing mechanisms for ensuring that interventions achieve their intended outcomes while maintaining high standards of quality accountability (Institute of Medicine, 2003). Quality assurance systems should include standardized protocols, training programs, and oversight mechanisms that ensure consistent implementation across different settings and providers. Performance measurement systems should include both process and outcome indicators that can track progress toward environmental health goals while identifying areas for improvement. Regular evaluation and quality improvement activities should be integrated into ongoing program operations, with mechanisms for incorporating feedback into program modifications. Transparency and public reporting of performance data can enhance accountability while building public support for environmental health programs (Olajide et al., 2023).

#### 4. Conclusion

The examination of conceptual frameworks linking pollution exposure, climate change, and public health outcomes reveals a complex and evolving landscape that requires sophisticated, integrated approaches to environmental health protection and disease prevention. This comprehensive review has demonstrated that environmental health challenges operate through multiple interconnected pathways that span direct toxicological effects, ecosystem-mediated impacts, and socially-mediated vulnerabilities that collectively shape population health outcomes across diverse contexts and scales. The evidence clearly indicates that traditional approaches focused on single pollutants or single exposure pathways are inadequate for addressing the complexity of real-world environmental health challenges, necessitating the development of more sophisticated frameworks that can capture cumulative exposures, mixture effects, and the dynamic interactions between environmental and social determinants of health.

Climate change has emerged as a fundamental driver that is reshaping the environmental health landscape, acting both as a direct determinant of health outcomes and as a threat multiplier that exacerbates existing environmental health risks while creating new exposure scenarios and vulnerability patterns. The climate-health linkages identified in this review operate across multiple temporal and spatial scales, from acute health effects during extreme weather events to gradual changes in disease transmission patterns and ecosystem health that unfold over decades. These relationships demonstrate the urgent need for environmental health frameworks that can address both immediate adaptations needs and long-term prevention strategies while recognizing the interconnected nature of climate, environment, and health systems. The evidence suggests that effective climate change adaptation strategies must be integrated with broader environmental health protection efforts to achieve optimal health outcomes while building resilience to future environmental challenges.

The role of social determinants in shaping environmental health outcomes represents a critical dimension that must be central to any comprehensive environmental health framework. This review has documented extensive evidence of environmental health disparities across racial, ethnic, and socioeconomic lines, with marginalized communities consistently facing disproportionate exposure environmental hazards while having limited access to resources for protection and recovery. These patterns reflect structural inequalities and systemic injustices that cannot be addressed through technical interventions alone, but require comprehensive approaches that address the root causes of environmental health disparities while building community capacity for environmental health advocacy and action. The concept of environmental justice provides essential grounding for frameworks that can promote equity while addressing environmental health challenges.

The challenges and barriers to environmental health integration identified in this review highlight the need for systemic changes in how environmental health is approached across institutions, sectors, and scales of action. Institutional fragmentation, limited capacity, funding constraints, and political barriers create significant obstacles to effective environmental health protection that cannot be overcome through isolated interventions. Instead, these challenges require coordinated efforts to build institutional capacity,

develop supportive policy environments, and create sustainable financing mechanisms that can support long-term environmental health improvement efforts. The evidence suggests that addressing these systemic barriers is essential for achieving meaningful progress in environmental health outcomes.

The best practices and recommendations presented in this review provide a roadmap for developing and implementing more effective environmental health frameworks that can address current challenges while building capacity for future environmental health protection. Systems-based approaches, community engagement strategies, integrated surveillance systems, and adaptive management frameworks represent core components of effective environmental health practice that have demonstrated success across diverse contexts. These approaches emphasize the importance of moving beyond traditional public health practice to embrace interdisciplinary, multi-sectoral, and community-centered strategies that can address the complexity of environmental health challenges while promoting equity and sustainability. The integration of technological innovations and data analytics represents an important opportunity for advancing environmental health practice, with new tools for exposure assessment, health surveillance, and intervention delivery creating possibilities for more precise, timely, and effective environmental responses. However, health implementation of these technologies must be guided by principles of equity and accessibility to ensure that technological advances do not exacerbate existing disparities or create new barriers to environmental health protection. The evidence suggests that technology should be viewed as a tool to support broader environmental health goals rather than as an end in itself.

The policy implications of this review are substantial, highlighting the need for comprehensive policy frameworks that can address environmental health across multiple sectors while promoting equity and sustainability. Health in All Policies approaches provide valuable frameworks for integrating health considerations into decision-making processes across sectors, while environmental justice principles provide essential guidance for ensuring that policy interventions address the needs of vulnerable populations. The evidence suggests that effective environmental health policy requires sustained political commitment, adequate resources, and mechanisms for community engagement and accountability.

The research implications of this review point to several important areas where additional investigation is needed to advance environmental health knowledge and practice. Priority areas include developing better methods for assessing cumulative exposures and mixture effects, understanding the mechanisms through which social determinants modify environmental health relationships, evaluating the effectiveness of different intervention strategies across diverse contexts, and examining the long-term health impacts of climate change and environmental degradation. The evidence also suggests the need for more interdisciplinary research approaches that can capture the complexity of environment-health relationships while producing actionable knowledge for policy and practice.

The practice implications of this review emphasize the need for significant changes in how environmental health is approached within public health and related fields. Environmental health practitioners need enhanced training and capacity to address the complexity of current environmental health challenges, while public health institutions need to develop greater capacity for interdisciplinary collaboration, community engagement, and systems thinking. The evidence suggests that effective environmental health practice requires moving beyond traditional approaches focused on individual exposures and health effects to embrace more comprehensive strategies that address the social, economic, and political determinants of environmental health outcomes.

Looking toward the future, several trends and developments are likely to shape the evolution of environmental health frameworks and practice. Climate change will continue to create new environmental health challenges while exacerbating existing risks, requiring adaptive frameworks that can respond to changing conditions and emerging threats. Technological advances will create new opportunities for environmental health assessment and intervention, while also potentially creating new risks that require proactive assessment and regulation. Urbanization, globalization, and demographic changes will alter exposure patterns and vulnerability profiles, requiring frameworks that can address these shifting patterns while promoting health equity.

The growing recognition of planetary health concepts represents an important development that may influence future environmental health frameworks by emphasizing the interconnections between human health and the health of natural systems. This perspective recognizes that human health is fundamentally dependent on healthy ecosystems and that environmental health strategies must consider the sustainability of natural systems that support human wellbeing. The planetary health framework may provide valuable guidance for developing environmental health approaches that promote both human health and environmental sustainability.

The importance of global cooperation and knowledge sharing in addressing environmental health challenges is likely to increase as environmental problems become increasingly transboundary and global in scope. Climate change, air pollution, chemical contamination, and other environmental health threats require coordinated responses across national boundaries, creating needs for international frameworks that can support collaborative action while respecting local contexts and priorities. The development of global environmental health governance mechanisms represents an important area for future development.

In conclusion, the conceptual frameworks linking pollution exposure, climate change, and public health outcomes represent a rapidly evolving area that requires continued innovation and adaptation to address emerging challenges while building on existing knowledge and successful practices. The evidence presented in this review demonstrates both the complexity of environmental health challenges and the potential for effective interventions that can protect and promote population health while addressing environmental degradation and social inequities. Achieving this potential will require sustained commitment, adequate resources, and collaborative approaches that can bridge traditional disciplinary and sectoral boundaries while centering equity and community engagement in environmental health efforts. The path forward requires recognition that environmental health is not simply a technical problem requiring technical solutions, but a complex challenge that reflects broader social, economic, and political systems that shape both

environmental conditions and health outcomes. Effective responses must address these systemic issues while building capacity for ongoing adaptation and improvement as conditions change and knowledge advances. The frameworks and approaches outlined in this review provide important guidance for this work, but their successful implementation will depend on the commitment and capacity of individuals, organizations, and communities working together to create healthier, more equitable, and more sustainable futures for all populations.

#### 5. References

- Adeleke O, Ajayi SAO. Transforming the healthcare revenue cycle with artificial intelligence in the USA; 2024
- 2. Afrihyia E, Umana AU, Appoh M, Frempong D, Akinboboye O, Okoli I, *et al.* Enhancing software reliability through automated testing strategies and frameworks in cross-platform digital application environments. J Front Multidiscip Res. 2022;3(2):517-31. doi:10.54660/.JFMR.2022.3.1.517-531
- 3. Agyeman J, Bullard RD, Evans B. Just sustainabilities: development in an unequal world. Cambridge: MIT Press; 2016.
- 4. Ajayi SAO, Akanji OO. Efficacy of mobile health apps in blood pressure control in USA. [place unknown: publisher unknown]; 2022.
- 5. Ajayi SAO, Akanji OO. Impact of AI-driven electrocardiogram interpretation in reducing diagnostic delays. [place unknown: publisher unknown]; 2023.
- 6. Akhamere GD. Behavioral indicators in credit analysis: predicting borrower default using non-financial behavioral data. Int J Manag Organ Res. 2022;1(1):258-66. doi:10.54660/JJMOR.2022.1.1.258-266
- 7. Akhamere GD. Fairness in credit risk modeling: evaluating bias and discrimination in AI-based credit decision systems. Int J Adv Multidiscip Res Stud. 2023;3(6):2061-70.
- 8. Akhamere GD. The impact of Central Bank Digital Currencies (CBDCs) on commercial bank credit creation and financial stability. Int J Adv Multidiscip Res Stud. 2023;3(6):2071-9.
- 9. Akinboboye IO, Okoli I, Frempong D, Afrihyia E, Omolayo O, Appoh M, *et al.* Applying predictive analytics in project planning to improve task estimation, resource allocation, and delivery accuracy. Int J Multidiscip Res Growth Eval. 2022;3(4):675-89. doi:10.54660/.IJMRGE.2022.3.4.675-689
- 10. Akinboboye O, Afrihyia E, Frempong D, Appoh M, Omolayo O, Umar MO, *et al.* A risk management framework for early defect detection and resolution in technology development projects. Int J Multidiscip Res Growth Eval. 2021;2(4):958-74. doi:10.54660/.IJMRGE.2021.2.4.958-974
- 11. Akintayo OD, Ifeanyi CN, Onunka O. Enhancing domestic peace through effective community-based ADR programs. Glob J Adv Res Rev. 2024;2(2):1-15.
- 12. Alonge EO, Eyo-Udo NL, Chibunna B, Ubanadu AID, Balogun ED, Ogunsola KO. Data-driven risk management in US financial institutions: a theoretical perspective on process optimization. Iconic Res Eng J. 2023;7(3):[page range unavailable].
- 13. Annan CA. Mineralogical and geochemical characterisation of monazite placers in the Neufchâteau

- Syncline (Belgium). [place unknown: publisher unknown]; 2021.
- 14. Appoh M, Frempong D, Akinboboye O, Okoli I, Afrihyia E, Umar MO, *et al.* Agile-based project management strategies for enhancing collaboration in cross-functional software development teams. J Front Multidiscip Res. 2022;3(2):49-64. doi:10.54660/.IJFMR.2022.3.2.49-64
- 15. Awe T. Cellular localization of iron-handling proteins required for magnetic orientation in C. elegans. [place unknown: publisher unknown]; 2021.
- 16. Ayumu MT, Ohakawa TC. Optimizing public-private partnerships (PPP) in affordable housing through fiscal accountability frameworks, Ghana in focus. IRE J. 2021;5(6):332-9.
- 17. Ayumu MT, Ohakawa TC. Real estate portfolio valuation techniques to unlock funding for affordable housing in Africa. Int J Multidiscip Res Growth Eval. 2022;3(1):967-72.
- 18. Ayumu MT, Ohakawa TC. Adaptive underutilizedal strategies: converting underutilized commercial properties into affordable housing. Int J Multidiscip Res Growth Eval. 2023;4(1):1200-6.
- 19. Ayumu MT, Ohakawa TC. Financial modeling innovations for affordable housing development in the U.S. Int J Adv Multidiscip Res Stud. 2024;4(6):1761-6.
- 20. Bai X, van der Leeuw S, O'Brien K, Berkhout F, Biermann F, Brondizio ES, *et al.* Plausible and desirable futures in the Anthropocene: a new research agenda. Glob Environ Change. 2016;39:351-62.
- 21. Barbier EB, Hacker SD, Kennedy C, Koch EW, Stier AC, Silliman BR. The value of estuarine and coastal ecosystem services. Ecol Monogr. 2008;78(3):235-53.
- 22. Basu R. High ambient temperature and mortality: a review of epidemiologic studies from 2001 to 2008. Environ Health. 2009;8:40.
- 23. Beck L, Frankel A. A conceptual model for continuing competence for environmental health practice. J Environ Health. 2004;66(9):22-8.
- 24. Ben-Shlomo Y, Kuh D. A life course approach to chronic disease epidemiology: conceptual models, empirical challenges and interdisciplinary perspectives. Int J Epidemiol. 2002;31(2):285-93.
- 25. Berry HL, Whitman A, Lyons RA, Tattersall R, Lavoie J. The case for systems thinking about climate change and mental health. Nat Clim Chang. 2018;8(4):282-90.
- 26. Brook RD, Rajagopalan S, Pope CA 3rd, Brook JR, Bhatnagar A, Diez-Roux AV, *et al.* Particulate matter air pollution and cardiovascular disease: an update to the scientific statement from the American Heart Association. Circulation. 2010;121(21):2331-78.
- 27. Brown P. Popular epidemiology and toxic waste contamination: lay and professional ways of knowing. J Health Soc Behav. 1992;33(3):267-81.
- 28. Brownson RC, Fielding JE, Maylahn CM. Evidence-based public health: a fundamental concept for public health practice. Annu Rev Public Health. 2009;30:175-201.
- 29. Brunekreef B, Holgate ST. Air pollution and health. Lancet. 2002;360(9341):1233-42.
- 30. Bullard RD, Johnson GS. Environmental justice: grassroots activism and its impact on public policy decision making. J Soc Issues. 2000;56(3):555-78.
- 31. Burke TA, Cascio WE, Costa DL, Deener K, Fontaine

- TD, Fulk FA, *et al.* Rethinking environmental health for the 21st century: recommendations of a national environmental health research strategy. Environ Health Perspect. 2017;125(11):117003.
- 32. Clayton S. Climate anxiety: psychological predispositions and climate change worry. Glob Environ Change. 2020;62:102263.
- 33. Cohen AJ, Brauer M, Burnett R, Anderson HR, Frostad J, Estep K, *et al.* Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. Lancet. 2017;389(10082):1907-18
- 34. Collins PH, Bilge S. Intersectionality. Hoboken: John Wiley & Sons; 2016.
- 35. Corburn J. Street science: community knowledge and environmental health justice. Cambridge: MIT Press; 2005.
- Cutter SL, Boruff BJ, Shirley WL. Social vulnerability to environmental hazards. Soc Sci Q. 2003;84(2):242-61.
- 37. Dannenberg AL, Bhatia R, Cole BL, Heaton SK, Feldman JD, Rutt CD. Use of health impact assessment in the US: 27 case studies, 1999-2007. Am J Prev Med. 2008;34(3):241-56.
- 38. Davies GK, Davies MLK, Adewusi E, Moneke K, Adeleke O, Mosaku LA, *et al.* AI-enhanced culturally sensitive public health messaging: a scoping review. E-Health Telecommun Syst Netw. 2024;13(4):45-66.
- Davies JC. Nanotechnology oversight: an agenda for the next administration. Environ Law Rep. 2009;39:10363-
- 40. Dogho MO. A literature review on arsenic in drinking water. [place unknown: publisher unknown]; 2021.
- 41. Dogho MO. Adapting solid oxide fuel cells to operate on landfill gas: methane passivation of Ni anode. Youngstown: Youngstown State University; 2023.
- 42. Dominici F, Peng RD, Barr CD, Bell ML. Protecting human health from air pollution: shifting from a single-pollutant to a multi-pollutant approach. Epidemiology. 2010;21(2):187-94.
- 43. Ebi KL, Semenza JC. Community-based adaptation to the health impacts of climate change. Am J Prev Med. 2008;35(5):501-7.
- 44. Ebi KL, Semenza JC, Rocklöv J. Current medical research funding and frameworks are insufficient to address the health risks of global environmental change. Environ Health. 2016;15:108.
- 45. Ebi KL, Woodruff R, von Hildebrand A, Corvalan C. Climate change, sustainable development, and health: moving forward by stepping back. Ecohealth. 2018;15(1):12-23.
- 46. Edwards QC, Smallwood S. Accessibility and comprehension of United States health insurance among international students: a gray area. [place unknown: publisher unknown]; 2023.
- 47. Ezeamii JC, Edwards QC, Omale J, Ezeamii PC, Idoko B, Ejembi EV. Risk beyond the pap: a review of key epidemiological studies on cervical cancer risk factors and populations at highest risk. World J Adv Res Rev. 2024;24(2):1402-20.
- 48. Fagbore OO, Ogeawuchi JC, Ilori O, Isibor NJ, Odetunde A, Adekunle BI. Building cross-functional collaboration models between compliance, risk, and

- business units in finance. [place unknown: publisher unknown]; 2024.
- 49. Finn S, O'Fallon L. The emergence of environmental health literacy—from its roots to its future potential. Environ Health Perspect. 2017;125(4):495-501.
- 50. Flanagan BE, Gregory EW, Hallisey EJ, Heitgerd JL, Lewis B. A social vulnerability index for disaster management. J Homel Secur Emerg Manag. 2011;8(1):3.
- 51. Folke C, Carpenter SR, Walker B, Scheffer M, Chapin T, Rockström J. Resilience thinking: integrating resilience, adaptability and transformability. Ecol Soc. 2010;15(4):20.
- 52. Frempong D, Akinboboye O, Okoli I, Afrihyia E, Umar MO, Umana AU, *et al.* Real-time analytics dashboards for decision-making using Tableau in public sector and business intelligence applications. J Front Multidiscip Res. 2022;3(2):65-80. doi:10.54660/.IJFMR.2022.3.2.65-80
- 53. Frempong D, Umana AU, Umar MO, Akinboboye O, Okoli I, Omolayo O. Multi-tool collaboration environments for effective stakeholder communication and sprint coordination in agile project teams. Int J Sci Res Comput Sci Eng Inf Technol. 2024;10(4):606-45. doi:10.32628/IJSRCSEIT
- 54. Frieden TR. A framework for public health action: the health impact pyramid. Am J Public Health. 2010;100(4):590-5.
- 55. Friel S, Marmot M, McMichael AJ, Kjellstrom T, Vågerö D. Global health equity and climate stabilisation: a common agenda. Lancet. 2008;372(9650):1677-83.
- 56. Frumkin H, Frank L, Jackson R. Urban sprawl and public health: designing, planning, and building for healthy communities. Washington: Island Press; 2004.
- 57. Gee GC, Payne-Sturges DC. Environmental health disparities: a framework integrating psychosocial and environmental concepts. Environ Health Perspect. 2004;112(17):1645-53.
- 58. Glasgow RE, Vinson C, Chambers D, Khoury MJ, Kaplan RM, Hunter C. National Institutes of Health approaches to dissemination and implementation science: current and future directions. Am J Public Health. 2012;102(7):1274-81.
- 59. Goldmann E, Galea S. Mental health consequences of disasters. Annu Rev Public Health. 2014;35:169-83.
- 60. Grandjean P, Bellanger M. Calculation of the disease burden associated with environmental chemical exposures: application of toxicological information in health economic estimation. Environ Health. 2017;16:123.
- 61. Hancock T. Population health promotion 2.0: an ecosocial approach to public health in the Anthropocene. Can J Public Health. 2015;106(4):e252-5.
- 62. Harris-Roxas B, Viliani F, Bond A, Cave B, Divall M, Furu P, *et al.* Health impact assessment: the state of the art. Impact Assess Proj Apprais. 2012;30(1):43-52.
- 63. Hayes K, Blashki G, Wiseman J, Burke S, Reifels L. Climate change and mental health risks in Australia: a prevention-focused adaptation approach. Int J Ment Health Syst. 2018;12:15.
- 64. Heindel JJ, Balbus J, Birnbaum L, Brune-Drisse MN, Grandjean P, Gray K, *et al.* Developmental origins of health and disease: integrating environmental influences. Endocrinology. 2015;156(10):3416-21.

- 65. Heinzerling L. Regulatory costs of mythic proportions. Yale Law J. 1998;107:1981-2070.
- 66. Hernández D. Understanding 'energy insecurity' and why it matters to health. Soc Sci Med. 2016;167:1-10.
- 67. Holling CS. Adaptive environmental assessment and management. Hoboken: John Wiley & Sons; 1978.
- 68. Hosking J, Campbell-Lendrum D. How well does climate change and human health research match the demands of policymakers? A scoping review. Environ Health Perspect. 2012;120(8):1076-82.
- 69. Howard G, Charles K, Pond K, Brookshaw A, Hossain R, Bartram J. Securing 2020 vision for 2030: climate change and ensuring resilience in water and sanitation services. J Water Clim Change. 2010;1(1):2-16.
- 70. Hunter PR. Climate change and waterborne and vector-borne disease. J Appl Microbiol. 2003;94 Suppl:37-46.
- 71. Ibidunni AS, William AAAA, Otokiti B. Adaptiveness of MSMEs during times of environmental disruption: exploratory study of capabilities-based insights from Nigeria. In: Innovation, entrepreneurship and the informal economy in Sub-Saharan Africa: a sustainable development agenda. Cham: Springer Nature Switzerland; 2024. p. 353-75.
- 72. Ilori O. Adopting zero trust security frameworks in financial and regulatory environments: a case study approach. [place unknown: publisher unknown]; 2022.
- 73. Ilori O, Kolawole TO, Olaboye JA. Ethical dilemmas in healthcare management: a comprehensive review. Int Med Sci Res J. 2024;4(6):703-25.
- 74. Institute of Medicine. The future of the public's health in the 21st century. Washington: National Academies Press; 2003.
- 75. Intergovernmental Panel on Climate Change. Climate change 2022: impacts, adaptation and vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge: Cambridge University Press; 2022.
- 76. Isa AK. Exploring digital therapeutics for mental health: AI-driven innovations in personalized treatment approaches. World J Adv Res Rev. 2024;24(3):10-30574.
- 77. Islam N, Winkel J. Climate change and social inequality. New York: UN Department of Economic and Social Affairs; 2017. Working Paper 152.
- 78. Israel BA, Eng E, Schulz AJ, Parker EA, editors. Methods for community-based participatory research for health. Hoboken: John Wiley & Sons; 2012.
- 79. Jacob DJ, Winner DA. Effect of climate change on air quality. Atmos Environ. 2009;43(1):51-63.
- 80. Jerrett M. Global geographies of health, exposure, and disease. In: A companion to health and medical geography. Hoboken: John Wiley & Sons; 2009. p. 10-32.
- 81. Jones AP. Indoor air quality and health. Atmos Environ. 1999;33(28):4535-64.
- 82. Kaggwa S, Onunka T, Uwaoma PU, Onunka O, Daraojimba AI, Eyo-Udo NL. Evaluating the efficacy of technology incubation centres in fostering entrepreneurship: case studies from the global south. Int J Manag Entrep Res. 2024;6(1):46-68.
- 83. Kasperson RE, Renn O, Slovic P, Brown HS, Emel J, Goble R, *et al.* The social amplification of risk: a conceptual framework. Risk Anal. 1988;8(2):177-87.

- 84. Kawachi I, Subramanian SV, Almeida-Filho N. A glossary for health inequalities. J Epidemiol Community Health. 2002;56(9):647-52.
- 85. Kortenkamp A, Backhaus T, Faust M. State of the art report on mixture toxicity. Brussels: European Commission; 2009.
- 86. Kriebel D, Tickner J, Epstein P, Lemons J, Levins R, Loechler EL, *et al.* The precautionary principle in environmental science. Environ Health Perspect. 2001;109(9):871-6.
- 87. Landrigan PJ, Fuller R, Acosta NJ, Adeyi O, Arnold R, Basu NN, *et al.* The Lancet Commission on pollution and health. Lancet. 2018;391(10119):462-512.
- 88. Lanphear BP, Hornung R, Khoury J, Yolton K, Baghurst P, Bellinger DC, *et al.* Low-level environmental lead exposure and children's intellectual function: an international pooled analysis. Environ Health Perspect. 2005;113(7):894-9.
- 89. Lasker RD, Weiss ES, Miller R. Partnership synergy: a practical framework for studying and strengthening the collaborative advantage. Milbank Q. 2001;79(2):179-205.
- Lichtveld M, Hodge J, St. Germain A, Arispe I, Sanderson LM. Lessons learned about emergency preparedness and response from the 2005 hurricanes. Disaster Med Public Health Prep. 2009;3(2 Suppl):S56-61.
- 91. Lowe A, Norris AC, Farris AJ, Babbage DR. Quantifying thematic saturation in qualitative data analysis. Field Methods. 2018;30(3):191-207.
- 92. Malin SA, Ryder SS. Developing deeply intersectional environmental justice scholarship. Environ Sociol. 2018;4(1):1-7.
- 93. Marmot M, Wilkinson R, editors. Social determinants of health. Oxford: Oxford University Press; 2006.
- 94. McMichael AJ, Montgomery H, Costello A. Health risks, present and future, from global climate change. BMJ. 2012;344:e1359.
- 95. Mendell MJ. Indoor residential chemical emissions as risk factors for respiratory and allergic effects in children: a review. Indoor Air. 2007;17(4):259-77.
- Millennium Ecosystem Assessment. Ecosystems and human well-being: synthesis. Washington: Island Press; 2005.
- 97. Minkler M, Wallerstein N, editors. Community-based participatory research for health: from process to outcomes. Hoboken: John Wiley & Sons; 2012.
- 98. Morello-Frosch R, Lopez R. The riskscape and the color line: examining the role of segregation in environmental health disparities. Environ Res. 2006;102(2):181-96.
- 99. Myers SS, Zanobetti A, Kloog I, Huybers P, Leakey AD, Bloom AJ, *et al.* Increasing CO2 threatens human nutrition. Nature. 2014;510(7503):139-42.
- 100.National Research Council. Science and decisions: advancing risk assessment. Washington: National Academies Press; 2009.
- 101.Nkwazema OC, Dokuchits EY, Ejairu ED. Integrated asset management for extreme disasters: assessing the 2023 flood events and resilience of critical infrastructures. [place unknown: publisher unknown]; [year unknown].
- 102.Norberg J, Cumming GS, editors. Complexity theory for a sustainable future. New York: Columbia University Press; 2008.

- 103.Northridge ME, Stover GN, Rosenthal JE, Sherard D. Environmental equity and health: understanding complexity and moving forward. Am J Public Health. 2003;93(2):209-14.
- 104. Nwankwo TC, Ejairu E, Awonuga KF, Usman FO, Nwankwo EE. Conceptualizing sustainable supply chain resilience: critical materials manufacturing in Africa as a catalyst for change. Int J Sci Res Arch. 2024;11(1):2427-37
- 105.O'Neill BC, Oppenheimer M, Warren R, Hallegatte S, Kopp RE, Pörtner HO, *et al.* IPCC reasons for concern regarding climate change risks. Nat Clim Chang. 2017;7(1):28-37.
- 106.Ogden NH, Lindsay LR. Effects of climate and climate change on vectors and vector-borne diseases: ticks are different. Trends Parasitol. 2016;32(8):646-56.
- 107.Okoli I, Akinboboye O, Frempong D, Omolayo O. Optimizing academic operations with spreadsheet-based forecasting tools and automated course planning systems. Int J Multidiscip Res Growth Eval. 2022;3(4):658-74. doi:10.54660/.IJMRGE.2022.3.4.658-674
- 108.Okoye CC, Ofodile OC, Tula ST, Nifise AOA, Falaiye T, Ejairu E, *et al.* Risk management in international supply chains: a review with USA and African cases. Magna Sci Adv Res Rev. 2024;10(1):256-64.
- 109.Oladeinde BH, Olaniyan MF, Muhibi MA, Uwaifo F, Richard O, Omabe NO, *et al.* Association between ABO and Rh blood groups and hepatitis B virus infection among young Nigerian adults. J Prev Med Hyg. 2022;63(1):E109.
- 110.Olajide JO, Otokiti BO, Nwani S, Ogunmokun AS, Adekunle BI, Fiemotongha JE. Designing cash flow governance models for public and private sector treasury operations. Int J Sci Res Civ Eng. 2023;7(6):45-54.
- 111. Olajide JO, Otokiti BO, Nwani S, Ogunmokun AS, Adekunle BI, Efekpogua J. Designing integrated financial governance systems for waste reduction and inventory optimization. [place unknown: publisher unknown]; 2020.
- 112.Olajide JO, Otokiti BO, Nwani S, Ogunmokun AS, Adekunle BI, Efekpogua J. Developing a financial analytics framework for end-to-end logistics and distribution cost control. [place unknown: publisher unknown]; 2020.
- 113.Olaniyan MF, Uwaifo F, Olaniyan TB. Antiinflammatory, viral replication suppression and hepatoprotective activities of bitter kola-lime juicehoney mixture in HBeAg seropositive patients. Matrix Sci Pharma. 2022;6(2):41-5.
- 114.Olulaja O, Afolabi O, Ajayi S. Bridging gaps in preventive healthcare: telehealth and digital innovations for rural communities. In: Illinois Minority Health Conference; 2024 Oct; Naperville, IL. Springfield: Illinois Department of Public Health; 2024.
- 115.Omolayo O, Akinboboye O, Frempong D, Umana AU, Umar MO. Defect detection strategies in agile teams: improving software quality through automation and collaborative workflows. Int J Sci Res Comput Sci Eng Inf Technol. 2023;9(5):519-55. doi:10.32628/IJSRCSEIT
- 116.Palinkas LA. Global climate change, population displacement, and public health: the global south meets the global north. Cham: Springer International

- Publishing; 2020.
- 117.Peckham TK, Baker D, Camp JE, Kaufman JD, Seixas NS. Creating a future for occupational health. Ann Work Expo Health. 2017;61(1):3-15.
- 118.Pellow DN. Total liberation: the power and promise of animal rights and the radical earth movement. Minneapolis: University of Minnesota Press; 2017.
- 119.Prüss-Ustün A, Wolf J, Corvalán C, Bos R, Neira M. Preventing disease through healthy environments: a global assessment of the burden of disease from environmental risks. Geneva: World Health Organization; 2016.
- 120.Pulido L. Rethinking environmental racism: white privilege and urban development in Southern California. Ann Assoc Am Geogr. 2000;90(1):12-40.
- 121. Quinn MM, Kriebel D. Occupational health and safety in the global economy: workplace hazards and inequality in a changing world. Int J Occup Environ Health. 2006;12(3):232-45.
- 122.Rajagopalan S, Al-Kindi SG, Brook RD. Air pollution and cardiovascular disease: JACC state-of-the-art review. J Am Coll Cardiol. 2018;72(17):2054-70.
- 123.Rechtschaffen C, Markell DL. Reinventing environmental enforcement and the state/federal relationship. Washington: Environmental Law Institute; 2003.
- 124.Reid CE, Brauer M, Johnston FH, Jerrett M, Balmes JR, Elliott CT. Critical review of health impacts of wildfire smoke exposure. Environ Health Perspect. 2016;124(9):1334-43.
- 125.Richardson SD, Ternes TA. Water analysis: emerging contaminants and current issues. Anal Chem. 2018;90(1):398-428.
- 126.Rider CV, Dinse GE, Umbach DM, Simmons JE, Hertzberg RC. A strategy to assess toxicological interactions among multiple chemical exposures. Toxicol Appl Pharmacol. 2018;354:234-44.
- 127.Rider CV, Furr JR, Wilson VS, Gray LE Jr. A mixture of seven antiandrogens induces reproductive malformations in rats. Int J Androl. 2010;33(2):443-62.
- 128.Rizwan AM, Dennis LY, Chunho LIU. A review on the generation, determination and mitigation of urban heat island. J Environ Sci. 2008;20(1):120-8.
- 129.Rothman KJ, Greenland S, Lash TL. Modern epidemiology. Philadelphia: Lippincott Williams & Wilkins; 2008.
- 130.Ruckelshaus M, McKenzie E, Tallis H, Guerry A, Daily G, Kareiva P, *et al.* Notes from the field: lessons learned from using ecosystem service approaches to inform realworld decisions. Ecol Econ. 2015;115:11-21.
- 131.Rudd R, Buttke D. Integrating environmental health into veterinary medical curricula. J Vet Med Educ. 2012;39(2):126-32.
- 132.Rudolph L, Caplan J, Ben-Moshe K, Dillon L. Health in all policies: a guide for state and local governments. Washington: American Public Health Association; 2013.
- 133.Rydin Y, Bleahu A, Davies M, Dávila JD, Friel S, De Grandis G, *et al.* Shaping cities for health: complexity and the planning of urban environments in the 21st century. Lancet. 2012;379(9831):2079-108.
- 134.Sampson RJ, Raudenbush SW, Earls F. Neighborhoods and violent crime: a multilevel study of collective efficacy. Science. 1997;277(5328):918-24.
- 135.Sandman PM. Hazard versus outrage in the public

- perception of risk. In: Effective risk communication. New York: Springer; 1989. p. 45-9.
- 136.Schecter A, Birnbaum L, Ryan JJ, Constable JD. Dioxins: an overview. Environ Res. 2006;101(3):419-28
- 137. Schillinger D, Grumbach K, Piette J, Wang F, Osmond D, Daher C, *et al.* Association of health literacy with diabetes outcomes. JAMA. 2002;288(4):475-82.
- 138. Schlosberg D, Collins LB. From environmental to climate justice: climate change and the discourse of environmental justice. Wiley Interdiscip Rev Clim Change. 2014;5(3):359-74.
- 139.Sejoro S, Chatterjee A, Ayo-Farai O, Edwards Q, Adhikari A. Impact of gaseous air pollutants and climate change on airborne pollen load in the coastal area of Charleston—a five-year study. In: APHA 2024 Annual Meeting and Expo; 2024 Oct; Minneapolis, MN. Washington: American Public Health Association; 2024.
- 140. Semenza JC, Menne B. Climate change and infectious diseases in Europe. Lancet Infect Dis. 2009;9(6):365-75.
- 141.Sexton K, Hattis D. Assessing cumulative health risks from exposure to environmental mixtures—three fundamental questions. Environ Health Perspect. 2007;115(5):825-32.
- 142.Sexton K, Adgate JL, Ramachandran G, Pratt GC, Mongin SJ, Stock TH, *et al.* Comparison of personal, indoor, and outdoor exposures to hazardous air pollutants in three urban communities. Environ Sci Technol. 2004;38(2):423-30.
- 143.Sexton K, Olden K, Johnson BL. "Environmental justice": the central role of research in establishing a credible scientific foundation for informed decision making. Toxicol Ind Health. 1993;9(5):685-727.
- 144. Sheffield PE, Landrigan PJ. Global climate change and children's health: threats and strategies for prevention. Environ Health Perspect. 2011;119(3):291-8.
- 145.Shendell DG, Alexander MS, McCarty FA, Atlantis E, Morawetz J. The long-term effects of air pollution on respiratory health: the adult outcomes of childhood exposure. Respir Med. 2005;99(10):1287-92.
- 146.Shendell DG, Prill R, Fisk WJ, Apte MG, Blake D, Faulkner D. Associations between classroom CO2 concentrations and student attendance in Washington and Idaho. Indoor Air. 2004;14(5):333-41.
- 147.Swartjes FA, editor. Dealing with contaminated sites: from theory towards practical application. Dordrecht: Springer Science & Business Media; 2011.
- 148. Teutsch SM, Thacker SB. Planning a public health surveillance system. Epidemiol Bull. 1995;16(1):1-6.
- 149. Thacker SB, Qualters JR, Lee LM. Public health surveillance in the United States: evolution and challenges. MMWR Suppl. 2012;61(3):3-9.
- 150.Umana AU, Afrihyia E, Appoh M, Frempong D, Akinboboye O, Okoli I, *et al.* Data-driven project monitoring: leveraging dashboards and KPIs to track performance in technology implementation projects. J Front Multidiscip Res. 2022;3(2):35-48. doi:10.54660/.IJFMR.2022.3.2.35-48
- 151.Uwaifo F, Uwaifo AO. Bridging the gap in alcohol use disorder treatment: integrating psychological, physical, and artificial intelligence interventions. Int J Appl Res Soc Sci. 2023;5(4):1-9.
- 152. Vermeulen R, Schymanski EL, Barabási AL, Miller GW. The exposome and health: where chemistry meets

- biology. Science. 2020;367(6476):392-6.
- 153. Vrijheid M. Health effects of residence near hazardous waste landfill sites: a review of epidemiologic literature. Environ Health Perspect. 2000;108 Suppl 1:101-12.
- 154. Vrijheid M. The exposome: a new paradigm to study the impact of environment on health. Thorax. 2014;69(9):876-8.
- 155.Watts N, Adger WN, Agnolucci P, Blackstock J, Byass P, Cai W, *et al.* Health and climate change: policy responses to protect public health. Lancet. 2015;386(10006):1861-914.
- 156.Watts N, Amann M, Arnell N, Ayeb-Karlsson S, Beagley J, Belesova K, *et al.* The 2020 report of The Lancet Countdown on health and climate change: responding to converging crises. Lancet. 2021;397(10269):129-70.
- 157. Wheeler T, von Braun J. Climate change impacts on global food security. Science. 2013;341(6145):508-13.
- 158. Whitmee S, Haines A, Beyrer C, Boltz F, Capon AG, de Souza Dias BF, *et al.* Safeguarding human health in the Anthropocene epoch: report of The Rockefeller Foundation—Lancet Commission on planetary health. Lancet. 2015;386(10007):1973-2028.
- 159. Wild CP. Complementing the genome with an "exposome": the outstanding challenge of environmental exposure measurement in molecular epidemiology. Cancer Epidemiol Biomarkers Prev. 2005;14(8):1847-50.
- 160. Williams DR, Mohammed SA. Discrimination and racial disparities in health: evidence and needed research. J Behav Med. 2009;32(1):20-47.
- 161. Wing S. Environmental justice, science, and public health. Environ Health Perspect. 2005;113(1):A54-5.
- 162. Woodcock J, Edwards P, Tonne C, Armstrong BG, Ashiru O, Banister D, *et al.* Public health benefits of strategies to reduce greenhouse-gas emissions: urban land transport. Lancet. 2009;374(9705):1930-43.
- 163. Woodruff TJ, Sutton P, Navigation Guide Work Group. An evidence-based medicine methodology to bridge the gap between clinical and environmental health sciences. Health Aff (Millwood). 2014;33(7):1176-84.
- 164.Zinsstag J, Schelling E, Waltner-Toews D, Tanner M. From "one medicine" to "one health" and systemic approaches to health and well-being. Prev Vet Med. 2011;101(3-4):148-56.